



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification <sup>7</sup> : <b>A23L 3/00</b></p>	<b>A2</b>	<p>(11) International Publication Number: <b>WO 00/48477</b></p> <p>(43) International Publication Date: 24 August 2000 (24.08.00)</p>												
<div style="display: flex; justify-content: space-between;"> <div style="width: 48%;"> <p>(21) International Application Number: <b>PCT/US00/03993</b></p> <p>(22) International Filing Date: 14 February 2000 (14.02.00)</p> <p>(30) Priority Data:</p> <table style="width: 100%; border: none;"> <tr> <td style="width: 30%;">09/253,482</td> <td style="width: 40%;">19 February 1999 (19.02.99)</td> <td style="width: 30%;">US</td> </tr> <tr> <td>Not furnished</td> <td>9 February 2000 (09.02.00)</td> <td>US</td> </tr> </table> <p>(71) Applicant (for all designated States except US): MORN- INGSTAR DIAGNOSTICS, INC. [US/US]; Suite 130, 1376 Lead Hill Boulevard, Roseville, CA 95661 (US).</p> <p>(72) Inventors; and</p> <p>(75) Inventors/Applicants (for US only): KEMP, Maurice, Clarence [US/US]; 4401 Woodthrush Drive, El Dorado Hills, CA 95762 (US). LALUM, Robert, Blaine, Lalum [US/US]; 6949 Greenback Lane #B, Citrus Heights, CA 95621 (US). XIE, Zhong, Wei [US/US]; 1141 Elderberry Circle, Folsom, CA 95630 (US). CUNHA, Michael, Anthony [US/US]; 355 Shadow Ridge Drive #3, Roseville, CA 95678 (US). CARPENTER, Robert, H. [US/US]; 1303 Pecan Street, Bastrop, TX 78602 (US). ZHANG, Shu [US/US]; 401 Rocky Point Drive #5, Roseville, CA 95678 (US). YAO, Yu [US/US]; 401 Rocky Point Drive #5, Roseville, CA 95678 (US). LEWIS, David, E. [US/US]; 816 3rd Avenue, Eau Claire, WI 54703 (US).</p> </div> <div style="width: 48%;"> <p>(74) Agent: CHWANG, T., Ling; Hitt Chwang &amp; Gaines, P.C., Suite 225, 275 West Campbell Road, Richardson, TX 75080 (US).</p> <p>(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p><b>Published</b> Without international search report and to be republished upon receipt of that report.</p> </div> </div>			09/253,482	19 February 1999 (19.02.99)	US	Not furnished	9 February 2000 (09.02.00)	US						
09/253,482	19 February 1999 (19.02.99)	US												
Not furnished	9 February 2000 (09.02.00)	US												
<p>(54) Title: <b>ACIDIC SOLUTION OF SPARINGLY-SOLUBLE GROUP IIA COMPLEXES</b></p> <p>(57) Abstract</p> <p>An acidic solution of sparingly-soluble Group IIA complexes ("AGIIS"), its preparation and its uses. The AGIIS can be prepared by mixing a mineral acid (such as sulfuric acid), and a Group IIA hydroxide (such as calcium hydroxide) or a Group IIA salt of a dibasic acid (such as calcium sulfate), or a mixture of the two Group IIA compounds, followed by removing the solid formed. The various uses include cleaning, food production, decontamination, bioremediation, agricultural application, medical application, and detoxification of substances.</p>														
<table border="1" style="margin-top: 10px;"> <caption>Data points for the graph</caption> <thead> <tr> <th>Base/Acid Ratio</th> <th>Normality</th> </tr> </thead> <tbody> <tr> <td>0.10</td> <td>29</td> </tr> <tr> <td>0.20</td> <td>14.5</td> </tr> <tr> <td>0.30</td> <td>7.25</td> </tr> <tr> <td>0.40</td> <td>3.6</td> </tr> <tr> <td>0.45</td> <td>0</td> </tr> </tbody> </table>			Base/Acid Ratio	Normality	0.10	29	0.20	14.5	0.30	7.25	0.40	3.6	0.45	0
Base/Acid Ratio	Normality													
0.10	29													
0.20	14.5													
0.30	7.25													
0.40	3.6													
0.45	0													

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LJ	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

ACIDIC SOLUTION OF SPARINGLY-SOLUBLE GROUP IIA COMPLEXES

-2-

## BACKGROUND

This application is a continuation-in-part of an application filed February 19, 1999, Serial No. 09/253,482, the entire content of which is hereby incorporated by reference.

This invention relates to an acidic solution of sparingly-soluble Group IIA- complexes ("AGIIS"), its preparations, and its uses.

10 In the late 80's and early 90's, researchers in Japan developed strong ionized water ("SIW") as disinfectants. The SIW was established as water with pH 2.7 or less, having an oxidation-reduction potential of 1,000 mv or more, and chlorine concentration of 0.8 ppm or more. The  
15 SIW is prepared by electrolysis of water.

Electrolysis of tap water has also been used to produce "strong acid water" and "strong alkali water" both of which were claimed to have antiseptic properties.

U.S. Patent Number 5,830,838 to Wurzbürger, et al.  
20 describes a solution for cleaning metal surfaces. The solution is prepared by mixing calcium hydroxide and potassium hydroxide with equivalent of sulfuric acid in water then passing the solution through a 10 micron filter. The resulting concentrate can be diluted depending on the  
25 degree of surface oxidation of the metal to be treated.

U.S. Patent Number 5,895,782 to Overton, et al. describes a solution for cleaning metal surfaces particularly non-ferrous alloys such as copper, brass and

-3-

high strength aluminum alloys. The solution is prepared by mixing  $\text{Ca}(\text{OH})_2$  and KOH with equivalent sulfuric acid in water then passing the solution through a 10 micron filter. The resulting concentrate can be used full strength or  
5 diluted depending on the degree of surface oxidation of the metal to be treated.

International Publication WO 94/09798 describes a pharmaceutical composition for treatment of disease, injury and other disorders. The pharmaceutical composition  
10 comprises a complex of a calcium-containing component and a sulfate-containing component in a pharmaceutically acceptable carrier. The reference teaches the isolation from natural materials, such as peat, the inorganic compositions. The inorganic preparations comprise an  
15 alkaline, aqueous or organic, or mixture thereof, extract of peat. Peat is extracted with aqueous solutions, organic solutions or water-miscible organic solvents at temperature from below room temperature up to the boiling point of the solvents. The preferred extracting solvents are those  
20 having a pH of at least 9. Biologically active constituents of fractionated peat preparations were identified as  $\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$  (gypsum),  $\text{CaSO}_4 \cdot \text{K}_2\text{SO}_4 \cdot \text{H}_2\text{O}$  (syngenite, also referred to as the double salt of gypsum) and  $\text{K}_3\text{Na}(\text{SO}_4)_2$  (apthitalite) by X-ray powder diffraction analysis. The  
25 reference also describes the synthesis of syngenite.

Chemists describe and measure the ability of a substance to donate protons  $[\text{H}^+]$  to a chemical reaction as the pKa of that substance where

-4-



Although a hydronium ion is usually represented by  $\text{H}^+$  or  $\text{H}_3\text{O}^+$ , but its true formula is not certain. The aggregate could be  $\text{H}_5\text{O}_2^+$ ,  $\text{H}_7\text{O}_3^+$ , or even  $\text{H}_9\text{O}_4^+$ .

5       Positively charged water has the ability to donate protons  $[\text{H}^+]$ . The donation of a proton is usually an intermediate step in any acid hydrolysis reaction. Acids are usually the chemical reagent used to donate protons in an aqueous solution. If the water could be the source of  
10 the  $[\text{H}^+]$ , then there would be fewer unwanted by-products (toxics) from the reactions and there would be less hazard associated with these products use.

A strong acid is used to neutralize and remove the lime, or quicklime, from the brick and mortar. A strong  
15 acid, such as hydrochloric acid, also known as muriatic acid, is also used to clean hard water spots on shower stalls, windows, glass, toilets, urinals, mirrors and other surfaces. Hydrochloric acid is used to de-scale water towers and heat exchangers and to adjust the pH of the  
20 waste water effluent. A full strength mineral acid, such as hydrochloric acid, is extremely corrosive to many substances, including metals. In addition, hydrochloric acid at a low pH of 0.5 or so will burn a human skin in seconds. The acid is also very harmful in that it emits  
25 fumes irritating to the mucous membrane. If left near other chemicals, like bleach, hydrochloric acid will interact with them, even through a typical plastic bottle.

-5-

It is thus desirable to be able to have a source of "acidity," or  $H_3O^+$ , without these unwanted disadvantages and be able to reduce environmental and safety hazards associated with acid hydrolysis. Preferably, this source of "acidity" should be able to prevent re-contamination following decontamination, not induce bacterial resistance, not alter the taste, color or smell of treated foodstuffs, not create any odor, effective in water in a wide range of temperatures, relatively free of danger when overdosed, can be neutralized after use, not carcinogenic or mutagenic, non-toxic, almost harmless in water and the environment, environmentally friendly, and can be stored for a long period of time without decomposition or turning into hazardous compound.

The control of microbial growth is necessary in many practical situations, and significant advances in agriculture, medicine and food science have been made through study of this area of microbiology. "Control of growth" means to prevent growth of microorganisms. This control is effected in one of two basic ways: (1) By killing microorganisms; or (2) by inhibiting the growth of microorganisms. Control of growth usually involves the use of physical or chemical agents which either kill or prevent the growth of microorganisms. Agents which kill cells are called "cidal" agents; agents which inhibit the growth of cells, but without killing them, are referred to as "static" agents. Thus the term "bactericidal" refers to killing bacteria and "bacteriostatic" refers to inhibiting the growth of bacterial cells. A "bactericide" kills

-6-

bacteria, a "fungicide" kills fungi. "Sterilization" is the complete destruction or elimination of all viable organisms in or on an object being sterilized. The object is either sterile or not, there are no degrees of sterilization. Sterilization procedures involve the use of heat, radiation or chemicals, or physical removal of microorganisms.

Microorganisms tend to colonize and replicate on different surfaces resulting in adherent heterogenous microbial accumulations termed "biofilms." Biofilms may form on surfaces of food substances, feed substances, and instrumentations. The microorganisms in the biofilms may include bacteria, fungi, viruses, and protozoans. Since food safety is a national priority, any product that can help by solving a multitude of problems associated with food production is desirable. Removal and control of biofilms which harbor dangerous microbial contamination is a sanitation goal that needs to be achieved. It is also desirable to be able to safely decontaminate water and nutrient by lowering pH to levels where contaminants would react and organisms cannot live.

As used herein, the term "nutriment" means something that nourishes, heals, or promotes growth and repair the natural wastage of organic life. Thus, food for a human and feed for an animal are all examples of nutriment. Other examples of nutriment include beverages, food additive, feed additive, beverage additive, food supplement, feed supplement, beverage supplement, seasoning, spices, flavoring agent, stuffing, food dressing, pharmaceutical,

-7-

biological product, and others. The nutriment can be of plant origin, animal origin, or synthetic. Current sanitizing, disinfectants and pesticides products on the market for these uses contain residues of chlorine, ammonia, organic iodine, metal salts and other deleterious residues. It is desirable to have a way that would preclude these residues by promoting acid hydrolysis without the presence of deleterious chemicals. Additionally, this method should generate few hazardous volatile gases. Importantly, it is highly desirable to have a composition that can control and the growth of, and kill, microorganisms and, at the same time, destroy the products, such as toxins, generated by, or associated with, the microorganisms.

#### SUMMARY

The present invention involves an acidic, or low pH, solution of sparingly-soluble Group IIA-complexes ("AGIIS"), its preparation, and its uses. One embodiment of the present invention pertains to highly acidic solution prepared by mixing or blending a mineral acid with a Group IIA hydroxide or a Group IIA salt of a dibasic acid, or a combination of the Group IIA hydroxide and Group IIA salt of a dibasic acid. Still other aspects of the present invention pertain to different methods to promote the safe, clean, and environmentally sensitive ways of chemical production, pharmaceutical production, cleaning, food production, decontamination, bioremediation, agricultural

-8-

application, medical application, and detoxification as well as decontamination of a wide variety of substances.

#### DESCRIPTION OF THE FIGURE

Figure 1 shows the relation of the desired final acid normality of AGIIS and the mole ratio of calcium hydroxide to sulfuric acid, given in moles of calcium hydroxide per mole of sulfuric acid.

#### DETAILED DESCRIPTION

One aspect of the present invention pertains to an acidic, or low pH, solution of sparingly-soluble Group IIA-complexes ("AGIIS"). The solution may have a suspension of very fine particles. The term "low pH" means the pH is below 7, in the acidic region. The AGIIS of the present invention with a certain acid normality does not have the same dehydrating behavior as sulfuric acid solution saturated with calcium sulfate having the same acid normality. In other words, the AGIIS of the present invention with a certain acid normality does not char sucrose as readily as does a saturated solution of calcium sulfate in sulfuric acid having the same normality. Further, the AGIIS is non-volatile at room temperature. It is less corrosive to a human skin than sulfuric acid saturated with calcium sulfate having the same acid normality. Not intending to be bound by the theory, it is believed that one embodiment of AGIIS comprises near-saturated, saturated, or super-saturated calcium, sulfate

-9-

anions or variations thereof, and/or complex ions containing calcium, sulfates, and/or variations thereof.

The term "complex," as used herein, denotes a composition wherein individual constituents are associated.

5 "Associated" means constituents are bound to one another either covalently or non-covalently, the latter as a result of hydrogen bonding or other inter-molecular forces. The constituents may be present in ionic, non-ionic, hydrated or other forms.

10 The acidic solution of sparingly-soluble Group IIA-complex salt ("AGIIS") can be prepared in several ways. Some of the methods involve the use of Group IA hydroxide but some of syntheses are devoid of the use of any added Group IA hydroxide, although it is possible that a small  
15 amount of Group IA metal may be present as "impurities." The preferred way of manufacturing AGIIS is not to add Group IA hydroxide to the mixture. As the phrase implies, AGIIS is highly acidic, ionic, with a pH of below about 2.

Wurzbarger, et al. in U.S. Patent 5,830,838 describes  
20 an acidic solution prepared by the "calcium-hydroxide/potassium-hydroxide method." The solution is produced by first adding two moles of concentrated sulfuric acid (93%) to 2 liters of de-ionized water. Separately, an aqueous solution of base is prepared by adding one mole of  
25 calcium hydroxide (hydrated lime) and two moles of potassium hydroxide to 20 liters of de-ionized water with stirring. The acid solution is then mixed with the base solution. The mixture is then filtered through a 10 micron filter to remove particles of calcium sulfate or

-10-

potassium sulfate of eleven microns or larger. The resulting concentrate can be used full strength or diluted with water depending on the metal surfaces to be treated. Sodium hydroxide may be used in place of potassium  
5 hydroxide. Hydrated calcium oxide may be used in place of calcium hydroxide. Another source of the base is calcium metal. In either case and as one embodiment of this application, the resultant solution is a highly acidic solution. This highly acidic solution can be diluted with  
10 water to adjust its pH to a desired higher value, i.e. less acidic.

Another way of preparing the acidic solution is by the "calcium-metal method" which involves reacting concentrated sulfuric acid with calcium metal followed by filtration.  
15 One mole of concentrated sulfuric acid was diluted with 40 moles of de-ionized water. Then, one mole of calcium metal turnings was slowly added with stirring into the solution of sulfuric acid. The stirring was continued until essentially all metal had dissolved. The resultant mixture  
20 was allowed to settle for about 5 to 6 hours before the supernatant was filtered through a 10 micron filter. The concentrate thus obtained had a pH value of about 0.5. This concentrate of hydronium ions was then diluted with de-ionized water to the desired pH value, such as pH of  
25 about 1 or about 1.8.

Then, there is the "calcium-hydride method" which involves reacting concentrated sulfuric acid and calcium hydride in water. One mole of concentrated sulfuric acid was diluted with 40 moles of de-ionized water. With

-11-

agitation, 1 mole of calcium hydride was slowly added to the solution of sulfuric acid. The agitation was continued until the calcium hydride has essentially all dissolved. After the dissolution, the mixture was then  
5 allowed to settle for about 5 to 6 hours, at that time the supernatant was filtered through a 10 micron filter. The concentrate thus obtained had a pH value of about 0.1 to about 0.2, and can be further diluted.

One product from the "calcium-metal method" or  
10 "calcium-hydride method" having a pH of from -0.2 to -0.3, and from 1.4 to 1.5 acid normality gave the following analyses: Ca, 763 ppm; SO<sub>4</sub>, 84633 ppm; Na, 4.76 ppm; K, 3.33 ppm; and Mg, 35.7 ppm.

The "calcium-metal method" and the "calcium-hydride  
15 method" have certain drawbacks. In each of these methods, thermal control is very difficult to achieve because of the large amount of heat generated when concentrated sulfuric acid is reacted with either calcium metal or calcium hydride. The difficulties in thermal control of the  
20 reactions cause the reactions to be difficult to reproduce and hard to control.

The preferred method of preparing AGIIS involves mixing a mineral acid with a Group IIA hydroxide, or with a Group IIA salt of a dibasic acid, or with a mixture of  
25 the two Group IIA materials. In the mixing, a salt of Group IIA is also formed. Preferably, the starting Group IIA material or materials selected will give rise to, and form, the Group IIA salt or salts that are sparingly soluble in water. The preferred mineral acid is sulfuric

-12-

acid, the preferred Group IIA hydroxide is calcium hydroxide, and the preferred Group IIA salt of a dibasic acid is calcium sulfate. Other examples of Group IIA salt include calcium oxide, calcium carbonate, and "calcium bicarbonate."

Thus, for example, AGIIS can be prepared by mixing or blending starting materials given in one of the following scheme with good reproducibility:

- (1)  $\text{H}_2\text{SO}_4$  and  $\text{Ca}(\text{OH})_2$ ;
- 10 (2)  $\text{H}_2\text{SO}_4$ ,  $\text{Ca}(\text{OH})_2$ , and  $\text{CaCO}_3$ ;
- (3)  $\text{H}_2\text{SO}_4$ ,  $\text{Ca}(\text{OH})_2$ ,  $\text{CaCO}_3$ , and  $\text{CO}_2$  (gas);
- (4)  $\text{H}_2\text{SO}_4$  and  $\text{CaCO}_3$ ;
- (5)  $\text{H}_2\text{SO}_4$ ,  $\text{CaCO}_3$ , and  $\text{Ca}(\text{OH})_2$ ;
- (6)  $\text{H}_2\text{SO}_4$ ,  $\text{CaCO}_3$ , and  $\text{CO}_2$  (gas);
- 15 (7)  $\text{H}_2\text{SO}_4$  and  $\text{CaSO}_4$ ;
- (8)  $\text{H}_2\text{SO}_4$ ,  $\text{Ca}(\text{OH})_2$ , and  $\text{CaSO}_4$ ;
- (9)  $\text{H}_2\text{SO}_4$ ,  $\text{CaSO}_4$ , and  $\text{CaCO}_3$ ;
- (10)  $\text{H}_2\text{SO}_4$ ,  $\text{CaSO}_4$ ,  $\text{CaCO}_3$ , and  $\text{Ca}(\text{OH})_2$ ;
- (11)  $\text{H}_2\text{SO}_4$ ,  $\text{CaSO}_4$ ,  $\text{CaCO}_3$ , and  $\text{CO}_2$  (gas); and
- 20 (12)  $\text{H}_2\text{SO}_4$ ,  $\text{CaSO}_4$ ,  $\text{CaCO}_3$ ,  $\text{CO}_2$  (gas), and  $\text{Ca}(\text{OH})_2$ .

Thus, preferably, AGIIS is prepared by mixing calcium hydroxide with concentrated sulfuric acid, with or without an optional Group IIA salt of a dibasic acid (such as calcium sulfate) added to the sulfuric acid. The optional calcium sulfate can be added to the concentrated sulfuric acid prior to the introduction of calcium hydroxide into the blending mixture. The addition of calcium sulfate to the concentrated sulfuric acid appears to reduce the amount

-13-

of calcium hydroxide needed for the preparation of AGIIS. Other optional reactants include calcium carbonate and gaseous carbon dioxide being bubbled into the mixture. Regardless of the use of any optional reactants, it was  
5 found that the use of calcium hydroxide is desirable.

One preferred method of preparing AGIIS can be described briefly as: Concentrated sulfuric acid is added to chilled water (8°-12°C) in the reaction vessel, then, with stirring, calcium sulfate is added to the acid in chilled  
10 water to give a mixture. Temperature control is paramount to this process. To this stirring mixture is then added a slurry of calcium hydroxide in water. The solid formed from the mixture is then removed. This method involves the use of sulfuric acid, calcium sulfate, and calcium  
15 hydroxide, and it has several unexpected advantages. Firstly, this reaction is not violent and is not exceedingly exothermic. Besides being easy to control and easy to reproduce, this reaction uses ingredients each of which has been reviewed by the U.S. Food and Drug  
20 Administration ("U.S. FDA") and determined to be "generally recognized as safe" ("GRAS"). As such, each of these ingredients can be added directly to food, subject, of course, to certain limitations. Under proper concentration, each of these ingredients can be used as  
25 processing aids and in food contact applications. Their use is limited only by product suitability and Good Manufacturing Practices ("GMP"). The AGIIS so prepared is thus safe for animal consumption, safe for processing aids, and safe in food contact applications. Further, the AGIIS

-14-

reduces biological contaminants in not only inhibiting the growth of, and killing, microorganisms but also destroying the toxins formed and generated by the microorganisms. The AGIIS formed can also preserve, or extend the shelf-life  
5 of, consumable products, be they plant, animal, pharmaceutical, or biological products. It also preserves or improves the organoleptic quality of a beverage, a plant product or an animal product. It also possesses certain healing and therapeutic properties.

10 The sulfuric acid used is usually 95-98% FCC Grade (about 35-37 N). The amount of concentrated sulfuric acid can range from about 0.05 M to about 18 M (about 0.1 N to about 36 N), preferably from about 1 M to about 5 M. It is application specific. The term "M" used denotes molar or  
15 moles per liter.

Normally, a slurry of finely ground calcium hydroxide suspended in water (about 50% of W/V) is the preferred way of introducing the calcium hydroxide, in increments, into the a stirring solution of sulfuric acid, with or without  
20 the presence of calcium sulfate. Ordinarily, the reaction is carried out below 40°C, preferably below room temperature, and more preferably below 10°C. The time to add calcium hydroxide can range from about 1 hour to about 4 hours. The agitation speed can vary from about 600 to  
25 about 700 rpm, or higher. After the mixing, the mixture is filtered through a 5 micron filter. The filtrate is then allowed to sit overnight and the fine sediment is removed by decantation.

-15-

The calcium hydroxide used is usually FCC Grade of about 98% purity. For every mole of concentrated acid, such as sulfuric acid, the amount, in mole, of calcium hydroxide used is application specific and ranges from  
5 about 0.1 to about 1.

The optional calcium carbonate is normally FCC Grade having a purity of about 98%. When used with calcium hydroxide as described above, for every mole of concentrated acid, such as sulfuric acid, the amount, in  
10 mole, of calcium carbonate ranges from about 0.001 to about 0.2, depending on the amount of calcium hydroxide used.

The optional carbon dioxide is usually bubbled into the slurry containing calcium hydroxide at a speed of from about 1 to about 3 pounds pressure. The carbon dioxide is  
15 bubbled into the slurry for a period of from about 1 to about 3 hours. The slurry is then added to the reaction vessel containing the concentrated sulfuric acid.

Another optional ingredient is calcium sulfate, a Group IIA salt of a dibasic acid. Normally, dihydrated  
20 calcium sulfate is used. As used in this application, the phrase "calcium sulfate," or the formula " $\text{CaSO}_4$ ," means either anhydrous or hydrated calcium sulfate. The purity of calcium sulfate (dihydrate) used is usually 95-98% FCC Grade. The amount of calcium sulfate, in moles per liter  
25 of concentrated sulfuric acid ranges from about 0.005 to about 0.15, preferably from about 0.007 to about 0.07, and more preferably from about 0.007 to about 0.04. It is application specific.

-16-

From experimental data, a slope was generated showing the ratio of calcium hydroxide to concentrated sulfuric needed for a desired final acid normality of AGIIS. See, Figure 1.

5        The slope in Figure 1 was created from two pairs of data points found by titrating a given amount of acid to a desired final acid normality. The accuracies were determined chemically. The final acid normality of the finished product ranges from about 1.2 to about 29. To  
10   produce one liter of 1.2 N AGIIS, it was found that for every mole of concentrated sulfuric acid, 0.45 moles of  $\text{Ca}(\text{OH})_2$  was required. To produce one liter of 27 N AGIIS, it was found that for every mole of concentrated sulfuric acid, 0.12 moles of  $\text{Ca}(\text{OH})_2$  was required. The data were  
15   then plotted onto a graph where the Y-axis represents final acid normality and the X-axis represents moles of  $\text{Ca}(\text{OH})_2$ /1 mole of concentrated sulfuric acid, where  $X_1 = 0.45$ ,  $X_2 = 0.12$ ,  $Y_1 = 1.2$ , and  $Y_2 = 27$ . The slope of the line was found by using the equation  $(Y_1 - Y_2)/(X_1 - X_2)$ , which was -  
20   78.18. The line can be represented by the equation  $Y = mX + b$ , where  $mX$  is the slope, and  $b$  is the Y intercept. The highest acid normality was 36.65, thus the equation is:

$$Y = -78.18X + 36.65$$

25        This slope is useful for the preparation of an AGIIS solution having a desired final acid normality.

Broadly, the method of preparing AGIIS having a desired final acid normality involves the steps given below. The calculations are based on a 1 liter of final

-17-

volume of AGIIS, the amounts of acid (concentrated sulfuric acid) and base (calcium hydroxide) are in moles, the ratio of base to acid is the number of moles of base (calcium hydroxide) for every mole of acid (concentrated sulfuric acid). The steps are:

(a) Determining the amount of mineral acid (such as concentrated sulfuric acid), in moles, needed to produce AGIIS having the desired final acid normality ("N") by using a relationship given by the following equation:

$$E_1 = (N/2) + (N/2 + B)$$

in which  $E_1$  is the amount of acid, in moles, required before correcting for purity, or purity adjustment; N is the desired final acid normality; and B is the mole ratio of the Group IIA hydroxide to the mineral acid needed to obtain the AGTIIS having N, and B is derived from a pre-plotted curve depicting the relationship of the mineral acid and the Group IIA hydroxide for a desired final N;

(b) making purity adjustment for the mineral acid used. The correction for the purity of the acid used is accomplished by the equation:

$$E_2 = E_1/C$$

in which  $E_2$  is the amount of acid, in moles, required after correcting for purity of the acid used, or purity adjustment;  $E_1$  is as defined above; and C is purity adjustment factor for the acid used. For concentrated sulfuric acid, the average acid strength is about 96.5%, and thus C is 0.965;

-18-

(c) determining the amount of water, in ml, that has to be added to the acid whose acid solution will then, after the reaction, give the desired final acid normality N. The relationship is as follows:

5 
$$G = J - E_2 - I$$

in which G is the amount of water required to be added to the mineral acid solution to get the desired final acid normality; J is the final volume of the aqueous mineral acid solution; I is the volume amount of Group IIA  
10 hydroxide needed (see, below); and  $E_2$  is as defined above;

(d) adding G to  $E_2$  to give the final aqueous solution of the mineral acid, in which both G and  $E_2$  are as defined above;

(e) determining the amount of base, (such as calcium  
15 hydroxide), in moles, needed for the reaction to produce AGIIS having the desired final acid normality N. For example, from the straight line in Figure 1, the mole ratio of  $\text{Ca}(\text{OH})_2$  to concentrated  $\text{H}_2\text{SO}_4$  to achieve a certain final acid normality can be determined.

20 the amount of the base, in moles, needed is:

$$F_1 = N/2 \times B$$

in which  $F_1$  is the amount of base, in moles, needed; and N and B are as defined above;

(f) the correction for the purity of the base used is  
25 accomplished by the equation:

$$F_2 = F_1/D$$

in which  $F_2$  is the amount of base, in moles, required after correcting for purity of the base used, or

-19-

purity adjustment; and D is purity adjustment factor for the base used.

The average purity of sodium hydroxide is about 98%, and, thus, D, in this case, is 0.98;

- 5 (g) determining the amount of water, in ml, needed to make the slurry of base. The relationship is as follows:

$$H = F_2 \times 1.5$$

in which H is the volume of water, in ml, needed to make the slurry of base which, in turn, will give AGIIS  
10 with the desired final acid normality N.  $F_2$  is as defined above. The H given is an approximation and should be adjusted to a desired final weight volume. Thus, for example, 50 g of base should be adjusted to a final volume of 100 ml because the slurry used is a 50 : 50 mixture of  
15 solid and water;

(h) determining the volume, in ml, of the base slurry or solution to be added to the acid solution to give AGIIS with the desired final acid normality N. The relationship can be expressed as:

20  $I = F_2 \times 2$

in which I is the volume, in ml, of the slurry or solution of base to be added to the acid solution; and  $F_2$  is as defined above;

- (i) adding H to  $F_2$  to give the final aqueous slurry or  
25 solution of the base, in which both H and  $F_2$  are as defined above;

(j) adding the final aqueous solution or slurry or the base of (i) to the final aqueous solution of mineral acid of (d);

-20-

(k) allowing the final aqueous solution or slurry of the base and the final aqueous solution of mineral acid (j) to react; and

(l) removing solid formed from the reaction of (k).

5 In the event that  $\text{CaSO}_4$  is used for the reaction by adding it to the solution of concentrated  $\text{H}_2\text{SO}_4$ , the amount of  $\text{CaSO}_4$ , in grams per liter of solution based on final volume, has the following relationship:

10	<u>Final AGIIS Acid Normality N</u>	<u>Amount of <math>\text{CaSO}_4</math> in g/l</u>
	1 - 5	5
	6-10	4
	11-15	3
	16-20	2
15	21-36	1

The AGIIS obtained could have an acid normality range of from about 0.05 to about 31; the pH of lower than 0; boiling point of from about 100 to about 106°C; freezing  
20 point of from about -8°C to about 0°C.

AGIIS obtained from using the reaction of  $\text{H}_2\text{SO}_4/\text{Ca}(\text{OH})_2/\text{CaSO}_4$  had the following analyses (average):

25 AGIIS With Final Acid Normality of 1.2 N, pH of  
-0.08

-21-

$\text{H}_3\text{O}^+$ , 2.22%; Ca, 602 ppm;  $\text{SO}_4$ , 73560 ppm; K, 1.36 ppb; impurities of 19.68 ppm, and neither Na nor Mg was detected.

AGIIS With Final Acid Normality of about 29 N, pH of  
5 about -1.46

$\text{H}_3\text{O}^+$ , 30.68%; Ca, 52.9 ppm;  $\text{SO}_4$ , 7356000 ppm; K, 38.02 ppb; and neither Na nor Mg was detected.

Besides concentrated sulfuric acid, other polyprotic acids, such as phosphoric acid, phosphorous acid, chloric acid, iodic acid, or others can be used.

Likewise, aqueous solutions of other alkalines or bases, such as Group IA hydroxide solution or slurry and Group IIA hydroxide solution or slurry can be used. Groups IA and IIA refer to the two Groups in the periodical table.

15 The use of Group IIA hydroxide is preferred. Preferably, the salts formed from using Group IIA hydroxides in the reaction are sparingly-soluble in water. It is also preferable to use only Group IIA hydroxide as the base without the addition of Group IA hydroxide.

20 After the reaction, the resultant concentrated acidic solution with a relatively low pH value, typically below pH 1, can then be diluted with de-ionized water to the desired pH value, such as pH of about 1 or about 1.8.

However, it is sometimes desirable not to prepare a very concentrated AGIIS solution and then dilute it serially to obtain the solution having the desired final acid normality. It is often desirable to prepare a

25

-22-

solution of AGIIS having a desired final pre-determined acid normality according to the method described in this application so that not much dilution of the product is required before use.

5       As discussed above, AGIIS has relatively less dehydrating properties (such as charring sucrose) as compared to the saturated solution of  $\text{CaSO}_4$  in the same concentration of  $\text{H}_2\text{SO}_4$ . Further, the stability and non-corrosive nature of the AGIIS of the present invention can  
10 be illustrated by the fact that a person can put his or her hand into this solution with a pH of less than 0.5 and, yet, his or her hand suffers no irritation, and no injury. If, on the other hand, one places his or her hand into a solution of sulfuric acid of less than pH 0.5, an  
15 irritation would occur within a relatively short span of time. A solution of 28 N of sulfuric acid saturated with calcium sulfate will cause chemical burn to a human skin after a few seconds of contact. In contrast, AGIIS solution of the same normality would not cause chemical  
20 burn to a human skin even after in contact for 5 minutes. The AGIIS of the present invention does not seem to be corrosive when being brought in contact with the environmental protective covering of plants (cuticle) and animals (skin). AGIIS is non-volatile at room temperature.  
25 Even as concentrated as 29 N, the AGIIS has no odor, does not give off fumes in the air, and is not irritating to a human nose when one smells this concentrated solution.

A "biological contaminant" is defined as a biological organism, or the product of biological organism, such as

-23-

toxin, or both, all of which contaminate the environment and useful products. This biological contaminant results in making the environment or product hazardous.

Biological contaminants, such as bacteria, fungi, 5 mold, mildew, spores, and viruses have potentially reactive substances in their cell wall/membranes; however, they hide in cells (viruses and some bacteria) and/or secrete biofilms (most bacteria, fungi, mold and mildew) to protect them from the environment.

10 Bacterial form or elaborate intracellular or extracellular toxins. Toxin is a noxious or poisonous substance that: (1) is an integral part of the bacteria; (2) is an extracellular product (exotoxin) of the bacteria; or (3) represents a combination of the two situations, 15 formed or elaborated during the metabolism and growth of bacteria. Toxins are, in general, relatively complex antigenic molecules and the chemical compositions are usually not known. The harmful effects of bacteria come not only from the bacteria themselves, but also from the 20 toxins produced by bacteria. Toxins produced by bacteria are just as, if not more, hazardous to the product than the bacteria themselves. Ordinary disinfectants, such as quaternary ammonium compounds, will kill bacteria but have no effect on bacterial toxins and endotoxins. In fact, 25 many disinfectants actually contribute to the endotoxins problems by causing their release from the bacteria. The bacterial toxins and endotoxins can cause serious adverse effects in human and animals. Endotoxins are the major cause of contamination in food products, in the production

-24-

of pharmaceuticals, medical devices, and other medical products. Thus, while "decontaminating" a product infested with bacteria, it is not enough to simply kill or reduce the number of bacteria. To get a safe and decontaminated product, the toxins and endotoxins of the bacteria must also be destroyed. Neither killing the microorganism alone nor destroying the toxins alone is enough in the real world. To be useful, when reducing biological contaminants in a nutriment or in an equipment, the growth of biological organisms must be controlled and reduced, and, at the same time, the product of biological organisms (such as toxins) must be removed and/or destroyed.

The outer covering, i.e. epidermis, of animals and cuticle of plants resist the growth and/or entry of the above microorganisms into the interior of the complex organism. One of the microbial growth prevention methods used by plants and animals is the maintenance of a surface pH or secretion of a coating that is not conducive to the attachment and propagation of micro-organisms. After a plant product is harvested or an animal product processed, these products lose the ability to resist the infestation of micro-organisms. By spraying the composition of the present invention plus defined additives on fruits, vegetables, and whole plants post harvest or washing or packing animal products in the composition, the growth and propagation of micro-organisms in these products can be reduced. If plant or animal products are packed in the composition an additional benefit is realized when the product is heated because the pH of the composition, and in

-25-

turn the product, goes down giving the added potential of the composition of destroying any micro-organisms, their toxins or other harmful substances.

The composition of the present invention was found to be a "preservative." The composition is not corrosive; however, it can create an environment where destructive micro-organisms cannot live and propagate, thus prolong the shelf-life of the product. The utility of this method of preservation is that additional chemicals do not have to be added to the food or other substance to be preserved because the inherent low pH of the mixture is preservative. Since preservative chemicals do not have to be added to the food substance, taste is improved and residues are avoided. Organoleptic testing of a number of freshly preserved and previously preserved food stuffs have revealed the addition of composition improves taste and eliminates preservative flavors. The term "organoleptic" means making an impression based upon senses of an organ or the whole organism. In another use, the composition was added to various food dressing, fresh juices and fermented beverages (wine). The resulting taste was unanimously judged better than the starting or control beverage. Use of the composition both as a preservative and taste enhancer for food and beverages will produce a safer and more desirable product. Additionally the composition can be added to biologics, pharmaceuticals and other preservative sensitive products to enhance their safety and extend shelf life. It can also be used as an ingredient to adjust product pH.

-26-

Conventional cleaning of biopharmaceutical and vaccine equipment is always problematical. Bioreactor vessels, where genetically altered yeast and bacteria produce biopharmaceutical products, are very sensitive to residues left during the cleaning process. The adduct or composition of the present invention is extremely useful in the primary cleaning of these vessels following production termination and for final cleaning and rinsing just prior to reestablishing the culture in the reactor vessel. The composition's ability to completely remove residues will insure the success of the culture and eliminate the possibility of contamination in the biopharmaceutical or vaccine product.

Another field of manufacturing where cleaning is critical is in the precision injection molding of plastic and composite materials for critical use parts in medical devices and other industrial products. The composition of the present invention can clean the injection molds quickly and efficiently between runs without damaging the molds or leaving residues which can cause defects in the product. Additionally, the composition could be used to remove excess materials from the parts and acid etch or clean parts prior to assembly and welding. The composition of the present invention is useful to clean the surface of non metallic parts to be chemically, heat or ultrasonically welded. If the device is wet packaged, i.e. suture material, then the composition can be used as a packaging preservative.

-27-

Agricultural applications for the composition of the present invention are of special interest. The ability to manipulate the pH of hydroponic plant production water will influence fruit production and disease control.

5 Synchronization of harvest and completeness of harvest can be aided by the composition. Olive, nut and some fruit trees are harvested by mechanical shaking. This shaking procedure must occur several times because the fruit and stem do not always ripen at the same time. Spraying the

10 tree with the composition prior to harvest activities can cause the stems and produce to mature rapidly. Only one or two shaking procedures will be required to completely harvest the produce, thus reducing harvest cost and damage to the trees.

15 Bacteria, fungus, yeast and molds can reduce plant yields or effect the quality of crops near, at, or post harvest. The composition of the present invention can be useful in preventing mold and mildew when crops in production are subjected to wet conditions. This is

20 especially true in corn, maize and other grain sorgham production. Grapes destined for raisin production are harvested and left to dry in the field on paper or cloth tarps between the vines. If wet weather persists the raisins will mold during the drying process resulting in an

25 unusable product. Spraying the composition on the grapes prior to harvest, dipping the clusters during harvest, treating the tarps, spraying the drying clusters, and washing the raisins prior to packing will result in raisins free of mold. The same methods can be used to assure

-28-

uniformity of grapes during wine making. The composition of the present invention can be used to control pH and adjust taste of wine and other fermented beverages.

The same use of the composition of the present invention can be made when storing grains. Mold, mildew and other fungal infestations of stored grains produce mycotoxins. These mycotoxins are very harmful to animals that consume contaminated grains. Mycotoxin intoxication results in organ damage, decreased production, or death.

10 Chemicals containing mercury and iodine are used to preserve planting seed, but there are no preservatives for grains destined for food or feed which do not leave harmful residues. Grains at harvest, during processing or in storage could be exposed to the composition, with or

15 without additives, to create an environment where these organisms would not grow on the grain or in the storage container.

Specific field applications for military use are numerous. The primary application is in the

20 decontamination of drinking water. Current methods for individual drinking water decontamination consist of placing iodine tablets into a canteen of water and waiting a period of time. If a small amount of the composition of the present invention is added to the water, time for

25 disinfection would be significantly reduced and there would be no need for iodine tablets. Additional applications for field living would include field waste decontamination, cooking liquid for food sources of questionable sanitary status, first aid irrigation solution for wounds and

-29-

decontamination, dilution and clean up of toxic or dangerous substance spills, and equipment cleaning and decontamination. This is especially important when food service under field conditions does not always allow for  
5 hot water cleaning of equipment.

The following examples are provided to further illustrate this invention and the manner in which it may be carried out. It will be understood, however, that the specific details given in the examples have been chosen for  
10 purposes of illustration only and not be construed as limiting the invention. Unless otherwise defined, the amount of each ingredient or component of the present invention is based on the weight percent of the final composition.

15

## Example 1

Preparation of 1.2-1.5 N AGIIS ( $H_2SO_4/Ca(OH)_2$ )

An amount of 1055 ml (19.2 moles, after purity adjustment and taking into account the amount of acid neutralized by base) of concentrated sulfuric acid (FCC  
20 Grade, 95-98% purity) was slowly added with stirring, to 16.868 L of RO/DI water in each of reaction flasks a, b, c, e, and f. The amount of water had been adjusted to allow for the volume of acid and the calcium hydroxide slurry. The mixture in each flask was mixed thoroughly. Each of  
25 the reaction flasks was chilled in an ice bath and the temperature of the mixture in the reaction flask was about 8-12°C. The mixture was continuously stirred at a rate of about 700 rpm.

-30-

Separately, a slurry was made by adding RO/DI water to 4 kg of calcium hydroxide (FCC Grace, 98% purity) making a final volume of 8 L. The mole ratio of calcium hydroxide to concentrated sulfuric acid was determined to be 0.45 to 1 from Figure 1. The slurry was a 50% (W/V) mixture of calcium hydroxide in water. The slurry was mixed well with a high-shear-force mixer until the slurry appeared uniform. The slurry was then chilled to about 8-12°C in an ice bath and continuously stirred at about 700 rpm.

10 To each of the reaction flasks was added 150 ml of the calcium hydroxide slurry every 20 minutes until 1.276 L (i.e. 638 g dry weight, 8.61 moles, of calcium hydroxide) of the slurry had been added to each reaction vessel. The addition was again accompanied by well mixing at about 700  
15 rpm.

After the completion of the addition of the calcium hydroxide to the reaction mixture in each reaction vessel, the mixture was filtered through a 5-micron filter.

The filtrate was allowed to sit for 12 hours, the  
20 clear solution was decanted to discard any precipitate formed. The resulting product was AGIIS having an acid normality of 1.2-1.5.

#### Example 2

##### Preparation of 2 N AGIIS ( $H_2SO_4/Ca(OH)_2/CaSO_4$ )

25 For the preparation of 1 L of 2 N AGIIS, an amount of 79.54 ml (1.44 moles, after purity adjustment and taking into account the amount of acid to be neutralized by base) of concentrated sulfuric acid (FCC Grade, 95-98% purity)

-31-

was slowly added, with stirring, to 853.93 ml of RO/DI water in a 2 L reaction flask. Five gram of calcium sulfate (FCC Grade, 95% purity) was then added slowly and with stirring to the reaction flask. The mixture was mixed  
5 thoroughly. At the point, the mixture would usually indicated an acid normality of 2.88. The reaction flask was chilled in an ice bath and the temperature of the mixture in the reaction flask was about 8-12°C. The mixture was continuously stirred at a rate of about 700  
10 rpm.

Separately, a slurry was made by adding 49.89 ml of RO/DI water to 33.26 g (0.44 mole, after purity adjustment) of calcium hydroxide (FCC Grace, 98% purity) making a final volume of 66.53 ml. The mole ratio of calcium hydroxide to  
15 concentrated sulfuric acid was determined to be 0.44 to 1 from Figure 1. The slurry was mixed well with a high-shear-force mixer until the slurry appeared uniform. The slurry was then chilled to about 8-12°C in an ice bath and continuous stirred at about 700 rpm.

20 The slurry was then slowly added over a period of 2-3 hours to the mixture, still chilled in an ice bath and being stirred at about 700 rpm.

After the completion of the addition of slurry to the mixture, the product was filtered through a 5-micron  
25 filter. It was normal to observe a 20% loss in volume of the mixture due to the retention of the solution by the salt and removal of the salt.

The filtrate was allow to sit for 12 hours, the clear solution was decanted to discard any precipitate formed.

-32-

The resulting product was AGIIS having an acid normality of 2.

### Example 3

#### Preparation of 12 N AGIIS ( $H_2SO_4/Ca(OH)_2/CaSO_4$ )

5 For the preparation of 1 L of 12 N AGIIS, an amount of 434.17 ml (7.86 moles, after purity adjustment and taking into account amount of acid neutralized by base) of concentrated sulfuric acid (FCC Grade, 95-98% purity) was slowly added, with stirring, to 284.60 ml of RO/DI water in  
10 a 2 L reaction flask. Three gram of calcium sulfate (FCC Grade, 95% purity) was then added slowly and with stirring to the reaction flask. The mixture was mixed thoroughly. The reaction flask was chilled in an ice bath and the temperature of the mixture in the reaction flask was about  
15 8-12°C. The mixture was continuously stirred at a rate of about 700 rpm.

Separately, a slurry was made by adding 210.92 ml of RO/DI water to 140.61 g (1.86 moles, after purity adjustment) of calcium hydroxide (FCC Grace, 98% purity)  
20 making a final volume of 281.23 ml. The mole ratio of calcium hydroxide to concentrated sulfuric acid was determined to be 0.31 from Figure 1. The slurry was mixed well with a high-shear-force mixer until the slurry appeared uniform. The slurry was then chilled to about 8-  
25 12°C in an ice bath and continuous stirred at about 700 rpm.

-33-

The slurry was then slowly added over a period of 2-3 hours to the mixture, still chilled in an ice bath and being stirred at about 700 rpm.

After the completion of the addition of slurry to the mixture, the product was filtered through a 5-micron filter. It was normal to observe a 20% loss in volume of the mixture due to the retention of the solution by the salt and removal of the salt.

The filtrate was allow to sit for 12 hours, the clear solution was decanted to discard any precipitate formed. The resulting product was AGIIS having an acid normality of 12.

#### Example 4

##### The Effects of AGIIS on Cold Sores

A forty-five year old white male discovered cold sores on his upper lip on day 1. He applied AGIIS, 4N, pH - 0.6, pH 1.8, to a cotton ball and "soaked" the sores for approximately one minute twice on day 1 and day 2. On day 3 he applied the AGIIS four times at various times throughout the day.

The slight pain that the cold sores caused was greatly reduced, almost immediately, upon applying the AGIIS to the sores. By the end of the third day of application the cold sores were virtually gone. Normally, it takes about seven days for the subject to heal cold sores using medication given to him by his physician.

AGIIS can be a useful treatment for cold sores due to herpes simplex.

- 34 -

AGIIS solution used in Examples 5 through Example 30 below was prepared by mixing concentrated sulfuric acid with either calcium hydride or calcium metal.

5 Example 5

## The Effects of AGIIS on Blade Shaver Cuts

A forty-five year old white male cut his face using a blade shaver in three locations. He applied the AGIIS, pH 1.8, with a "soaked" cotton ball directly to the cuts.

10       The cuts stopped bleeding within twenty seconds and  
the pain stopped almost immediately.

AGIIS can be useful as a cutaneous coagulant.

### Example 6

## Decontamination of Portable Water

15            Portable water contained non coliform organisms. AGIIS, pH 1.8, was added to this water to bring the pH to 2.0. There was no growth when the water was cultured, and the water could be consumed without adverse effects.

### Example 7

## 20 The Effects of AGIIS on Plaque and Bacteria

A forty-five year old white male with orthodontic appliances rinsed his mouth and teeth with AGIIS for 37 days. He used approximately 10 mL of AGIIS, pH 1.8, one to two times daily. He rinsed in the morning and sometimes  
25 prior to going to bed. He continued to brush his teeth twice per day and used an OTC mouthwash following the brushing.

-35-

He noticed that the surface of his teeth was not coated with a film as he had experienced prior to using AGIIS. He stated his mouth seemed to remain fresher for a longer period of time. He also noticed that his teeth  
5 seemed to be whiter and brighter. He received a dental cleaning on day 37. The hygienist performed a series of tests to evaluate the general condition of this teeth. The hygienist applied a dye to his teeth that allowed the hygienist to see plaque and/or bacteria that were present  
10 on his teeth. The hygienist used a computer with a video camera to view and record the condition of his teeth. The result showed that the top two thirds of his teeth showed virtually no plaque and no bacteria. The bottom one third that touches the gum area showed minor amounts of plaque  
15 and bacteria. The gums were determined to be in excellent condition. The hygienist suggested that the subject should wash with the AGIIS at least as often as he uses mouthwash and concentrate on bathing the gum area. The hygienist will continue to follow the progress. The hygienist also  
20 used a chemical and an ultraviolet light to determine if the AGIIS was removing tooth enamel. The study indicated that the AGIIS was not removing enamel.

AGIIS appears to help remove plaque and bacteria from the subjects teeth and mouth, whiten the teeth and kept his  
25 mouth fresher for a longer period of time without apparent removal of tooth enamel.

#### Example 8

##### Effect of AGIIS on a Tumor

-36-

A 50 year old man with multiple epidermoid cyst was topically treated with pH 1 AGIIS. Two tumor sites were selected and treated; however, there was no effect after 3 days. Then 0.1 mL of pH 1 AGIIS was injected intratumor  
5 via a 27 gauge needle and a tuberculin syringe. Within 24 hours the mass was gone and only a small scab where the mass was attached to the skin remained. There were no adverse effects and only slight stinging accompanied the injection. The scab at the tumor sight was gone in 7 days.

10

## Example 9

Effect of AGIIS on the Tissues of a Heparinized Dog

A 15 kg male beagle dog was scheduled for liver harvest to provide primary canine hepatocytes for toxicology tissue culture screening. The dog was prepared  
15 by having food withdrawn 24 hours prior to study. The dog was anesthetized by 2 mL of sodium pentothal and heparinized by injecting 5 mL, 1000 units/mL, heparin IV. Liver harvest was completed and various organs and skin incisions were exposed to an aqueous solution of a AGIIS  
20 having a pH value of 1. There were no adverse effects on the tissues exposed to the AGIIS. Heparinized blood placed in contact with AGIIS became brown and granular in color and consistency. There was no effect on clotting time in the heparinized dog.

25

## Example 10

Effect of AGIIS on Surgical/Wounds in a Rabbit

-37-

A male rabbit was anesthetized with 3 mL Ketamine IM, and his abdomen was shaved. Both sides of the abdomen were numbered with a permanent blue marker in the following sequence: 1, 2, 3, 4, 5, C where 1 = pH 1, 2 = pH 2, 3 = pH 3, 4 = water for irrigation ("WFI"), 5 = air control and C = clotting time control. The pH 1, 2, and 3 designate aqueous solutions of AGIIS having pH of 1, 2, and 3, respectively, or abbreviated as "pH 1 AGIIS treated" or "pH 1 treated," etc. Six incisions, 1 cm wide, were made at 2 different times. Various fluids corresponding to the labeled incision were introduced into the corresponding wound and the results observed for at least 20 minutes. Clotting times were determined by capillary tube fibrin method and found to be normal. Air control wounds clotted in about 2.5 minutes. Water for irrigation treated wounds appeared to have an extended clotting time of about 3 to 4 minutes duration. Wounds treated with an aqueous solution of AGIIS with pH 3 were not significantly different from WFI treated wounds. Wounds treated with an aqueous solution of AGIIS with pH 2 clotted in less than 2 minutes. Wounds treated with an aqueous solution of AGIIS with pH 1 clotted within 30 seconds and the clots assumed a dark brown halo around the periphery of the wound. By 5 minutes this wound was completely dry while all other wounds continued to ooze serum/lymph. All wounds were observed at 10 & 20 minutes. There were no differences noted at these observation times between the controls, WFI and pH 3 AGIIS treated wounds. At 20 minutes, the pH 2 treated wound was moist, but had contracted 10mm as

-38-

measured side to side. The pH 1 treated wound was dry with brown pigmented clots around the periphery of the wound. This wound had contracted 25mm and the subcutaneous tissues were light brown in color. This pigment was supposed to be  
5 hemosiderin which is the iron precipitant from the blood cells coming in contact with the AGIIS. Also of interest were the blood clots, while being brown on the outside were red and normal in appearance on the inside. All wounds were sutured with a 3-0 Vicryl ® mattress suture. Skin  
10 apposition was easier in the pH 1 AGIIS treated wound as the skin edges seemed to stick together as the suture was tied.

On the following day the incisions were examined and photographed. The incision treated with pH 1 AGIIS was  
15 more inflamed than the other incisions; however, it was completely closed. Only a small amount of pulling on the other incisions caused them to open, but equal and even increased tugging did not open the pH 1 AGIIS treated incisions. This finding was not expected. The tissue  
20 junction was dry and adhered. The rabbit was examined without anesthesia and did not display undue discomfort. There did not appear to be any effect on the synthetic Vicryl ® suture material.

#### Example 11

##### 25 Effect of AGIIS on the Ophthalmic Tissues of a Rabbit

The pH 1 and pH 2 AGIIS material was placed in the left and right eye respectively of a New Zealand white

-39-

rabbit. At the 10 minute observation, there appeared to be more redness in both eyes than normal; however, the rabbit did not and was not experiencing discomfort. At the 20 minute observation, there continued to be an increase in redness, but the eyes appeared normal. At the 1 hour observation, the eyes were slightly redder than normal, but the rabbit was not tearing or in discomfort. The rabbit was returned to his cage.

The above mentioned New Zealand white rabbit was examined approximately 24 hours post treatment. The eyes were examined and appeared normal. There was no evidence of corneal ulceration, opacity, or tearing.

#### Example 12

##### Effect and Use of AGIIS During a Surgical Procedure A 47

15 pound mixed female bull dog was presented for ovariohysterectomy. The dog was anesthetized with 10 mL (50mg/mL) pentabarbital sodium and intubated. The incision site was prepared with an alcohol and Betadine scrub. The incision was made with a #10 steel surgical blade. Large blood vessels were controlled with hemostats. An aqueous solution of AGIIS having pH of 1 was dropped via syringe onto small bleeding cutaneous vessels. While the hemorrhage was not stopped immediately, the tissues surrounding the vessels contracted exposing the bleeding vessels and facilitated their mechanical clamping. Very small vessels clotted immediately, as seen in the rabbit, and tissue fluid seepage into the surgical field was controlled. The ovaries and uterine horns were removed.

-40-

Two to four drops of the aqueous solution of AGIIS (pH = 1) were placed on the surgical stumps of the uterus and ovarian pedicles. Tissue color changed to slightly brown in tint, but there were no other tissue effects. The pH 1 AGIIS did not seem to have any effect on the peritoneal or serrosal surfaces of the abdominal organs. The skin edges of the incision were treated with pH 1 AGIIS prior to dog's closure. The closure with 2-0 Vicryl ® was routine. The dog was examined 24 hours later and the recovery and incision appeared normal. There were no adverse effects seen at the skin closure edges and the wound was sealed. The skin closure had a cosmetic appearance. Use of the pH 1 AGIIS did not appear to have any adverse effects on the surgically exposed tissues. It appeared to be effective in controlling hemorrhage in vessels less than 1 mm in outside diameter and lymphatics. Additionally, the AGIIS product rapidly removed blood from the surgical instruments.

#### Example 13

##### Investigation of pH 1.4 AGIIS to Remove Endotoxins from Glass Surfaces

Glass tubes were coated with BSA and autoclaved. Tube contents were removed and culture media along with E. coli 0157:H7 organisms were placed in the tubes. After incubation the tubes were autoclaved and the cycle was repeated in order to coat the tubes with endotoxins.

Tubes were divided into two groups: Group 1 tubes were filled with endotoxin free LAL water. Group II tubes was filled with pH 1.4 AGIIS solution. All tubes were then

-41-

boiled for 20 minutes. After boiling endotoxin free LAL water was introduced into each tube and the tubes were vortexed vigorously. The contents of each tube were assayed for endotoxins using an LAL Test Kit.

5        Testament with the pH 1.4 AGIIS solution decreased the associated endotoxin level from 22.66 EU/mL to undetectable levels ( $<0.03$  EU/mL). Treatment with LAL reagent water only did not reduce the endotoxin level associated with the glass tubes.

10 Example 14

Investigation of pH 1.4 AGIIS to Remove Endotoxins from  
Plastic Medical Devices

Plastic test tubes were coated with endotoxins by repeated culture with *E. coli* O157:H7 suspended in a beef suspension and autoclaving after each cycle. Tubes were divided into two groups. Group I tubes; boiled with endotoxin free LAL water. Group II tubes room temperature. Group III tubes; boiled with a pH 1.4 AGIIS solution. Treatment with the pH 1.4 AGIIS solution decreased the tube associated endotoxin level from ~45 EU/mL to undetectable levels (<0.03 EU/mL) or an ~256 fold reduction.

### Example 15

## Investigation of pH 1.4 AGIIS to Remove Endotoxins from Stainless Steel Surfaces

25           Stainless steel slabs (SSS) were coated with  
endotoxins by repeated culture with E. coli 0157:H7 and  
autoclaving after each cycle. SSS were divided into two

-42-

groups: Group 1 slabs were boiled with endotoxin LAL water. Group II coupons were boiled with a pH 1.4 AGIIS solution.

Treatment with the pH 1.4 AGIIS solution decreased the SSS associated endotoxin level from 4 EU/mL to undetectable levels (<0.03EU/mL). Treatment with LAL reagent water did not reduce the endotoxin level associated with the SSS.

#### Example 16

##### Anti-toxin Effect of AGIIS Treatment

10 An equal volume of a pH 0.5 solution of AGIIS was added to an E. coli O157:H7 culture. The resultant pH was ~1.0. The culture was then titrated back to ~pH 7.0 with 5N NaOH. The untreated and treated cultures were checked for Shiga Like Toxin II using Morningstar Diagnostic, Inc. SLT  
15 II test. The untreated culture was positive for SLT-II whereas the AGIIS treated culture was negative for SLT-II.

To show that we did not simply destroy all antigens, material from the untreated and AGIIS treated culture were checked for O157 antigens. Both the treated and AGIIS  
20 treated cultures were positive for O157 antigens. Therefore, the treatment with AGIIS either inactivated the toxin by destroying or dissociating the toxin to a non-antigenic form.

#### Example 17

25 Study to Determine if Different pH Solutions of AGIIS have Distinct Effects on the Oxidation of Bananas

-43-

Bananas were peeled and immersed in AGIIS solutions having a pH of 1.2, 1.4, 1.6, 1.8 or 2.0, respectively, for 5 min.

Oxidation of banana pieces was noticeable depressed by treatment with AGIIS solutions having a pH ranging from 1.2-1.6. After 24 hr bananas pieces treated with pH 1.2 and 1.4 AGIIS were for the most part free of oxidation. Thus low pH AGIIS is more effective at preventing oxidation of banana fruit pieces.

10

## Example 18

Study of pH 1.2 AGIIS in Prevention of Oxidation of Apples

Apples were cut in half and immersed in a pH 1.2 solution of AGIIS or in water. After treatment apple halves were removed and incubated at ambient room temperature. At 15 four hours post-treatment apple halves treated with the AGIIS solution were white while the water treated apple halves were brown due to oxidation. The differences were still apparent 24 hr later.

## Example 19

20 Study of pH 0.56 AGIIS in Removal of Oxidation from Brass Metal

Brass items were bathed in AGIIS solution and hard to remove oxidation was removed by scrubbing with stainless steel pads. Oxidation that accumulated over a twenty-year 25 period was removed with minimal effort.

## Example 20

-44-

Study of pH 0.56-AGIIS Solution in Decreasing the pH of a Sulfuric Acid Solution

Sulfuric acid was diluted to a pH 2.3 using deionized water (~700 mL). AGIIS solution added in 1 mL aliquots. pH went down in increments from 2.3 to 1.56. Therefore, a pH 0.56 solution of AGIIS could be used to increase the acidity of a sulfuric acid solution.

Example 21

Study to Determine the Concentration of a pH 0.45 AGIIS Solution

AGIIS (50 mL) was placed in an erlenmeyer and KOH or NaOH of known concentration (usually 1 N NaOH) was added to determine the "acidic" concentration of the AGIIS. Titration gave a value of 1.84 N. When base was added, the pH decreased from 0.45 to 0.35 and then increased steadily until neutrality was reached suggesting the dissociation of hydronium complexes in the presence of base to yield additional hydronium ions.

20

Example 22

Study to Assess the Effect of the Addition of AGIIS on the Organoleptic Properties of Wines

Cups were filled with 30 mL of wine. One hundred (100) microtiters of AGIIS (pH 0.3), were added to half of the cups and, 100 microliters of deionized water, were

-45-

added to the other half. A blinded panel of tasters was asked to taste the wine.

Changes in the organoleptic properties were noted. In particular, all tasters agreed the wine supplemented with  
5 AGIIS was less bitter. Color and pH of the wine were unchanged.

#### Example 23

##### Study to Determine the Effect of AGIIS on Concrete and Tile Surfaces

10 AGIIS applied at an ambient and elevated temperature to concrete removed grime and left the concrete between the stones whiter. The heated AGIIS was more effective than the ambient temperature AGIIS.

AGIIS applied to algae coated concrete killed and  
15 removed the algae.

Calcium carbonate deposits on swimming pool tiles were dissolved when AGIIS was applied.

AGIIS seems to be an effective agent for cleaning concrete surfaces without the corrosive effects of muriatic  
20 acid.

#### Example 24

##### Study to Determine if AGIIS Binds to Bran

Four 100-mL cups were filled with wheat bran. Two of the cups were filled with a pH 0.8 AGIIS solution while the  
25 remaining cups were filled with deionized water. The bran was allowed to rehydrate for 1 hr and all cups were then

-46-

placed in a -84°C freezer. Frozen cups were then placed in a lyophilizer for 24 hr.

After lyophilization the contents of each cup were removed and transferred to a 500-mL beaker. One hundred and fifty mL of pH 7 deionized water was added to each beaker and the freeze-dried bran was allowed to rehydrate.

Bran treated with AGIIS readily rehydrated and/or dissolved. Whereas the water treated bran had to be physically broken up before it dissolved.

10 When all samples were rehydrated, the pH of each sample was determined. The average pH of the bran treated with water was 5.8 whereas the pH of the AGIIS treated bran was 2.84. Thus AGIIS treatment lowered the pH of the treated bran and changed the rehydration characteristics of  
15 the bran.

#### Example 25

##### Effect of AGIIS on Oxidation of Avocados

Avocados were peeled and sliced into pieces.  
20 Individual pieces were immersed in AGIIS solutions having a pH of 1.2, 1.4, 1.6, 1.8 or 2.0, respectively, for 10 min. After incubation at ambient room temperature on an open shelf for 8 hrs, strong oxidation of pieces treated with pH 1.4-2.0 was evident. However, pieces treated with  
25 a pH 1.2 solution of AGIIS were free of oxidation and looked freshly cut.

#### Example 26

-47-

Study of the Effect on the Organoleptic Properties of Ketchup by Addition of AGIIS

Eighty milliliters of ketchup were placed in 100-mL cups. Five mL of deionized water was added to half of the  
5 cups. Five mL of AGIIS (~pH 0.5) was added to the other cups.

The cup contents were thoroughly mixed and a blinded panel of taste testers was asked to give their opinions and selection as to taste.

10 The AGIIS treated ketchup retained a thick consistency and the color stayed an intense red. Moreover, it was also determined that the taste was enhanced. The water treated ketchup lost consistency, color diminished and taste was judged not as good.

15

Example 27

Study of the Effect of AGIIS on a Plant Source of Pharmaceuticals

Freshly harvested aloe Vera leaf was dissected to  
20 expose the mucilaginous gel in the center of the leaf. Two sections were treated with AGIIS pH 2, and placed in an observation dish. Two other sections were treated with water and placed in an identical observation dish. After 10 minutes at room temperature, the water-treated aloe gel  
25 was discolored and appeared brown. The AGIIS-treated aloe gel retained its fresh-cut appearance. After 20 minutes at room temperature, the differences were even more pronounced. The water-treated gel began to liquify, while the AGIIS-treated gel retained its integrity. After four

-48-

hours at room temperature, the differences were even much more pronounced, and the AGIIS-treated gel still appeared freshly cut.

#### Example 28

##### 5 Effect of AGIIS on Contaminated Water

Bacteria present in 500 mL of tap water were concentrated by centrifugation at 5000 x g for 20 min. Another 500 mL of tap water was titrated to pH 2 using AGIIS solution of pH 0.5. Bacteria in the treated tap  
10 water was concentrated by centrifugation at 5000 x g for 20 min. Bacteria from each were suspended using 1.5 mL of the AGIIS or tap water and plated to determine the number of viable bacteria in each sample. Treatment with a pH 2 solution of AGIIS reduced the level of viable organism in  
15 the water.

#### Example 29

##### Effect of AGIIS on Street Puddle Water

Water was collected from a puddle at the corner in front of a laboratory building. It was determined that the  
20 pH of the water was 7.4. Water was mixed 1:1 with a pH 2 AGIIS solution or sterile saline and treated at ambient room temperature. Following treatment, an aliquot of the AGIIS- and saline- treated water was serially diluted and plated to determine the number of viable organisms. AGIIS  
25 treatment effectively decreased the number of viable organism relative to the control of saline.

-49-

## Example 30

Effect of AGIIS on Level of Viable Microbes on a Lettuce Head

5       Lettuce leafs were stripped from lettuce heads and placed in two groups. Group I lettuce leafs were treated with a pH 2 solution of AGIIS for 3 min and then stomached in sterile saline. Group II leafs were treated with saline for 3 min then stomached. An aliquot from each group was  
10       serially diluted and each dilution was plated to determine the number of viable organisms present following treatment. The number of viable organisms associated with a pH 2 solution of AGIIS was decreased compared to that of the control.

15

## Example 31

Effect of AGIIS On Hydrolysis of Chicken Feed

AGIIS was found to convert complex carbohydrates in chicken feed to monosaccharides which were much easier to digest than the complex carbohydrates in the stomach. The  
20       chicken feed was obtained from a commercial broiler producer. This grower ration contained 26% protein and was yellow corn based. The chicken feed was digested with 2 N of AGIIS at a temperature of 85° C for different length of time. AGIIS was prepared by  $H_2SO_4/Ca(OH)_2/CaSO_4$  method.  
25       Modified Fehling's solution method was employed to determine the amount of reducing sugar produced during the reaction. Controls using de-ionized water were performed in parallel. From the result given below, it can be seen

-50-

that chicken treated with AGIIS had higher amount of reducing sugar which is easier than complex carbohydrate for chicken to digest.

5	Sample Weight (g)	Reaction Time (hour)	Amount of Reducing Sugar (%)	
			AGIIS	Control
	15	1	2.96	0.2
	20	1	3.13	0
	25	1	4.85	0.1
	30	1	4.96	0.2
10	40	1	6.5	0.16
	40	2	8.1	0
	40	3	10.9	0
	40	4	14.2	0.33
	40	5	15.5	0.35
15	50	1	6.2	0.26

### Example 32

#### Charring of Sucrose by Various Agents

Sulfuric acid having a concentration of 19 N or higher will char or "dehydrate" sucrose. This reaction was visible and could be used as a measurement parameter. Results with sulfuric acid of less than 19 N was harder to

-51-

interpret due to the extended duration of the reaction. Roughly, the charring reaction could be divided into three stages.

The first stage was the initial color change. This usually occurred within the first two minutes of the reaction at room temperature. The first stage was characterized by color change in sucrose, i.e. the white color of sucrose turned into light yellow. Most acidic reagents used in this experiment would turn the color of sucrose into light yellow within the first two minutes of contact.

The second stage was the blackening of the sucrose.

The third stage was the charring or complete "burning" of sucrose. At this stage, heat was generated and vapor was given off. The reaction could be violent and mildly explosive depending on the concentration of the acid.

Given below is a table summarizing the results from comparative charring experiments of solutions of: (1) AGIIS; (2)  $H_2SO_4$ ; and (3)  $H_2SO_4 \cdot CaSO_4$ . The solution of AGIIS was prepared by the reaction of calcium hydroxide with sulfuric acid having added calcium sulfate therein. Solution (3), i.e.  $H_2SO_4 \cdot CaSO_4$ , was a solution of sulfuric acid saturated with calcium sulfate. The data were compiled from experiments carried out at room temperature.

- 52 -

	Solution	Initial Change	Time to Blacken	Time to Char
5	5 N AGIIS	No change	No change	Not Detected
	5 N H <sub>2</sub> SO <sub>4</sub> *CaSO <sub>4</sub>	No change	No change	Not Detected
	5 N H <sub>2</sub> SO <sub>4</sub>	No change	No change	Not Detected
	10 N AGIIS	>24 Hours	>24 Hours	Not Detected
	10 N H <sub>2</sub> SO <sub>4</sub> *CaSO <sub>4</sub>	>24 Hours	>24 Hours	Not Detected
	10 N H <sub>2</sub> SO <sub>4</sub>	>24 Hours	>24 Hours	Not Detected
10	19 N AGIIS	>20 min	~ Hour	Not Detected
	19 N H <sub>2</sub> SO <sub>4</sub> *CaSO <sub>4</sub>	<2 min	<1 Hour	Not Detected
	19 N H <sub>2</sub> SO <sub>4</sub>	40 sec	25 min	Not Detected
	27 N AGIIS	2 min	<10 min	Not Detected
15	27 N H <sub>2</sub> SO <sub>4</sub> *CaSO <sub>4</sub>	<2 min	<6 min	>10 min
	27 N H <sub>2</sub> SO <sub>4</sub>	Instant	<1 min	>10 min
	28 N AGIIS	<2 min	<10 min	Not Detected
	28 N H <sub>2</sub> SO <sub>4</sub> *CaSO <sub>4</sub>	<1 min	<5 min	<10 min
20	28 N H <sub>2</sub> SO <sub>4</sub>	Instant	<1 min	<10 min
	29 N AGIIS	1 min	<8 min	Not Detected
	29 N H <sub>2</sub> SO <sub>4</sub> *CaSO <sub>4</sub>	Instant	<5 min	<8 min
	29 N H <sub>2</sub> SO <sub>4</sub>	Instant	<1 min	<6 min

-53-

AGIIS, if prepared correctly, would cause the color of sucrose to remain yellow and only slowly darken over the next 7 or 8 minutes. AGIIS having an acid normality of between 27 and 29 N, if prepared incorrectly, will darken the color of sucrose in less than about 5 minutes. Further, the charring of the sucrose by properly prepared AGIIS, even at acid normality of 29 N, was not be detected more than 24 hours later at room temperature.

In contrast, as shown in the Table, either sulfuric acid or sulfuric acid saturated with calcium sulfate, under same acid normality, will char sucrose much more rapidly than AGIIS at room temperature.

### Example 33

#### Non-Volatility and Non-Corrosiveness of AGIIS

AGIIS prepared was non-volatile at room temperature. Even as concentrated as 29 N, the AGIIS had no odor, did not give off fumes in the air, and was not irritating to a human nose when one smelled the concentrated solution. When concentrated AGIIS was diluted with water, very little heat was given off, while dilution of concentrated sulfuric acid with water gave off a large amount of heat, i.e. very exothermic.

A human skin would get very hot upon contacting a solution of 28 N of sulfuric acid saturated with calcium sulfate. The solution was irritating to the skin within a few minutes, and chemical burn will follow. Sulfuric acid, 28 N, would chemically burn a human skin within less than one minute.

-54-

In contrast, upon contacting a human skin, a solution of AGIIS having an acid normality of 28 N, would cause only a mildly warm sensation. There was no irritating effects and the solution did not cause chemical burn even after about 5 minutes at room temperature on the skin.

**WHAT IS CLAIMED IS:**

1. A prepared nutriment comprising:
  - 2 a nutriment material; and
  - 3 an acidic sparingly-soluble Group IIA complex
  - 4 ("AGIIS").
2. The prepared nutriment of claim 1, wherein the
  - 2 AGIIS is isolated from a mixture comprising a mineral acid
  - 3 and a Group IIA hydroxide, or a Group IIA salt of a dibasic
  - 4 acid, or a mixture of the two.
3. The prepared nutriment of claim 2, wherein the
  - 2 Group IIA hydroxide is calcium hydroxide, the mineral acid
  - 3 is sulfuric acid, and the Group IIA salt of a dibasic acid
  - 4 is calcium sulfate.
4. The prepared nutriment of claim 3, wherein the
  - 2 AGIIS having a certain acid normality is less effective in
  - 3 charring sucrose and less corrosive to an animal skin than
  - 4 a saturated solution of calcium sulfate in sulfuric acid

-55-

5 having the same acid normality, and wherein the AGIIS is  
6 non-volatile at room temperature and pressure.

5. The prepared nutriment of claim 1, wherein the  
2 AGIIS, based on the total weight of the prepared nutriment,  
3 ranges from about 0.01 % to about 99.99 %.

6. The prepared nutriment of claim 1, wherein the  
2 nutriment material is food, feed, drink, food supplement,  
3 feed supplement, drink supplement, food dressing,  
4 pharmaceutical, biological product, seasoning, spices,  
5 flavoring agent, or stuffing.

7. A prepared nutriment comprising:  
2 a nutriment material; and  
3 AGIIS prepared by mixing calcium hydroxide and  
4 sulfuric acid with or without the addition of calcium  
5 sulfate.

8. The prepared nutriment of claim 7, wherein the  
2 sulfuric acid contains a predetermined amount of calcium  
3 sulfate.

9. The prepared nutriment of claim 7, wherein the  
2 AGIIS having a certain acid normality is less effective  
3 in charring sucrose and less corrosive to an animal skin  
4 than a saturated solution of calcium sulfate in sulfuric  
5 acid having the same acid normality, and wherein the  
6 AGIIS is non-volatile at room temperature and pressure.

-56-

10. The prepared nutriment of claim 7, wherein for  
2 every mole of sulfuric acid used, the amount of calcium  
3 hydroxide ranges from about 0.1 mole to about 0.5 mole.  
4

11. The prepared nutriment of claim 7, wherein the  
2 nutriment material is food, feed, drink, food supplement,  
3 feed supplement, drink supplement, food dressing,  
4 pharmaceutical, biological product, seasoning, spices,  
5 flavoring agent, or stuffing.

12. A method for manufacturing a prepared nutriment  
2 comprising:  
3 contacting AGIIS with a nutriment material.

13. A method for manufacturing a prepared nutriment  
2 comprising:  
3 contacting AGIIS with a carrier to give a  
4 constituted carrier; and  
5 blending the constituted carrier with a nutriment  
6 material.

14. A method for destroying organic odor in an  
2 environment, comprising:  
3 spraying the environment with AGIIS.

15. A method for preserving or improving  
2 organoleptic quality of a beverage, a plant product or an

-57-

3 animal product, comprising:           contacting the  
4 beverage, the plant product or the animal product with  
5 AGIIS.

16.     The method of claim 15, wherein the AGIIS is  
2 prepared by mixing calcium hydroxide with sulfuric acid  
3 with or without calcium sulfate added thereto, and the  
4 AGIIS having a certain acid normality is less effective  
5 in charring sucrose and less corrosive to an animal skin  
6 than a saturated solution of calcium sulfate in sulfuric  
7 acid having the same acid normality, and wherein the  
8 AGIIS is non-volatile at room temperature and pressure.

17.     A method for decreasing pH of AGIIS, the method  
2 comprising:  
3         heating the AGIIS.

18.     The method of claim 17, wherein the AGIIS is  
2 blended into food, feed, drink, food supplement, feed  
3 supplement, drink supplement, food dressing,  
4 pharmaceutical, biological product, seasoning, spices,  
5 flavoring agent, or stuffing.

19.     A method for reducing biological contaminants  
2 in a nutriment comprising:  
3         contacting the nutriment with AGIIS.

20.     The method of claim 19, wherein the AGIIS is  
2 prepared by mixing calcium hydroxide with sulfuric acid

-58-

3 with or without calcium sulfate added thereto, and the  
4 AGIIS having a certain acid normality is less effective  
5 in charring sucrose and less corrosive to an animal skin  
6 than a saturated solution of calcium sulfate in sulfuric  
7 acid having the same acid normality, and wherein the  
8 AGIIS is non-volatile at room temperature and pressure.

21. The method of claim 20, where the nutriment is  
2 a fresh fruit, a fruit product, a vegetable produce, a  
3 vegetable product, a meat, a meat product, a fish, a fish  
4 product, a food dressing, or a drink.

22. A method for reducing the pH of a nutriment  
2 comprising:  
3 contacting the nutriment with AGIIS.

23. The method of claim 22, wherein the AGIIS is  
2 prepared by mixing calcium hydroxide with sulfuric acid  
3 with or without calcium sulfate added thereto, and the  
4 AGIIS having a certain acid normality is less effective  
5 in charring sucrose and less corrosive to an animal skin  
6 than a saturated solution of calcium sulfate in sulfuric  
7 acid having the same acid normality, and wherein the  
8 AGIIS is non-volatile at room temperature and pressure.

24. A method for reducing biological contaminants  
2 in an equipment comprising:  
3 contacting the equipment with AGIIS.

-59-

25. The method of claim 23, wherein the AGIIS is  
2 prepared by mixing calcium hydroxide with sulfuric acid  
3 with or without calcium sulfate added thereto, and the  
4 AGIIS having a certain acid normality is less effective  
5 in charring sucrose and less corrosive to an animal skin  
6 than a saturated solution of calcium sulfate in sulfuric  
7 acid having the same acid normality, and wherein the  
8 AGIIS is non-volatile at room temperature and pressure.

26. The method of claim 24, wherein the equipment  
2 is a food-processing equipment, feed-processing  
3 equipment, drink-processing equipment, pharmaceutical  
4 equipment, construction equipment or micro-electronic  
5 equipment.

27. A method for preserving a consumable product,  
2 comprising:  
3 contacting the consumable product with AGIIS.

28. The method of claim 27, wherein the AGIIS is  
2 prepared by mixing calcium hydroxide with sulfuric acid  
3 with or without calcium sulfate added thereto, and the  
4 AGIIS having a certain acid normality is less effective  
5 in charring sucrose and less corrosive to an animal skin  
6 than a saturated solution of calcium sulfate in sulfuric  
7 acid having the same acid normality, and wherein the  
8 AGIIS is non-volatile at room temperature and pressure.

-60-

29. The method of claim 27, wherein the consumable  
2 product is a plant product, an animal product, a  
3 pharmaceutical product, a biological product, or a  
4 medical device product.

30. A method for reducing the quantity of a  
2 biological toxin in a medium, comprising:  
3 contacting the medium with AGIIS.

31. The method of claim 30, wherein the AGIIS is  
2 prepared by mixing calcium hydroxide with sulfuric acid  
3 with or without calcium sulfate added thereto, and the  
4 AGIIS having a certain acid normality is less effective  
5 in charring sucrose and less corrosive to an animal skin  
6 than a saturated solution of calcium sulfate in sulfuric  
7 acid having the same acid normality, and wherein the  
8 AGIIS is non-volatile at room temperature and pressure.

32. The method of claim 30, wherein the medium is a  
2 food, a feed, a pharmaceutical, an equipment, a packaging  
3 material, a drink, a biological product, water, or soil.

33. The method of claim 30, wherein the toxin is an  
2 animal toxin, a bacterial toxin, a botulinus toxin, a  
3 cholera toxin, a streptococcus erythrogenic toxin, a  
4 dinoflagellate toxin, a diphtheria toxin, an erythrogenic  
5 toxin, an extracellular toxin, a fatigue toxin, an  
6 intracellular toxin, a scarlet fever erythrogenic toxin,  
7 or a Tunnickliff toxin.

-61-

34. The method of claim 30, wherein the toxin  
2 comprises an endotoxin.

35. The method of claim 35, wherein the toxin  
2 comprises a mycotoxin.

36. A method for enhancing the bioavailability of a  
2 nutrient in a nutriment, comprising:  
3 adding to the nutriment AGIIS.

37. The method of claim 36, wherein the AGIIS is  
2 prepared by mixing calcium hydroxide with sulfuric acid  
3 with or without calcium sulfate added thereto, and the  
4 AGIIS having a certain acid normality is less effective  
5 in charring sucrose and less corrosive to an animal skin  
6 than a saturated solution of calcium sulfate in sulfuric  
7 acid having the same acid normality, and wherein the  
8 AGIIS is non-volatile at room temperature and pressure.

38. The method of claim 36, wherein the nutrient is  
2 a carbohydrate, a protein, an enzyme, or an acid-stable  
3 vitamin.

39. A method for incorporating AGIIS into a dry  
2 nutriment, comprising:  
3 adding AGIIS to a suitable carrier to give a  
4 premixed product, and

-62-

5       blending the premixed product with the dry  
6   nutriment.

40.     The method of claim 39, wherein the AGIIS is  
2   prepared by mixing calcium hydroxide with sulfuric acid  
3   with or without calcium sulfate added thereto, and the  
4   AGIIS having a certain acid normality is less effective  
5   in charring sucrose and less corrosive to an animal skin  
6   than a saturated solution of calcium sulfate in sulfuric  
7   acid having the same acid normality, and wherein the  
8   AGIIS is non-volatile at room temperature and pressure.

41.     The method of claim 39, wherein the suitable  
2   carrier is a methylcellulose, a psyllium, bran, rice hull  
3   or corn gluten.

42.     A method for treating a cutaneous anomaly on an  
2   animal, comprising:  
3       treating the cutaneous anomaly with AGIIS.

43.     The method of claim 42, wherein the AGIIS is  
2   prepared by mixing calcium hydroxide with sulfuric acid  
3   with or without calcium sulfate added thereto, and the  
4   AGIIS having a certain acid normality is less effective  
5   in charring sucrose and less corrosive to an animal skin  
6   than a saturated solution of calcium sulfate in sulfuric  
7   acid having the same acid normality, and wherein the  
8   AGIIS is non-volatile at room temperature and pressure.

-63-

44. The method of claim 42, wherein the cutaneous  
2 anomaly is a wound or a burn.

45. The method of claim 44, wherein the wound is a  
2 mechanical wound, a spontaneous ulceration, a dermatitis,  
3 or an eruption.

46. The method of claim 44, wherein the burn is a  
2 chemical burn or a thermal burn.

47. A method for inducing clotting of blood in a  
2 bleeding tissue in an animal, comprising:  
3 contacting the bleeding tissue with an AGIIS.

48. The method of claim 47, wherein the AGIIS is  
2 prepared by mixing calcium hydroxide with sulfuric acid  
3 with or without calcium sulfate added thereto, and the  
4 AGIIS having a certain acid normality is less effective  
5 in charring sucrose and less corrosive to an animal skin  
6 than a saturated solution of calcium sulfate in sulfuric  
7 acid having the same acid normality, and wherein the  
8 AGIIS is non-volatile at room temperature and pressure.

49. The method of claim 47, wherein the bleeding  
2 tissue is an external organ, an internal organ, a  
3 connective tissue, or a nerve tissue.

50. A method for enhancing the adhesion of a first  
2 tissue to a second tissue, comprising:

-64-

3           contacting AGIIS with the first tissue or both the  
4 first tissue and the second tissue; and  
5           joining the first tissue with the second tissue.

51.       The method of claim 50, wherein the AGIIS is  
2 prepared by mixing calcium hydroxide with sulfuric acid  
3 with or without calcium sulfate added thereto, and the  
4 AGIIS having a certain acid normality is less effective  
5 in charring sucrose and less corrosive to an animal skin  
6 than a saturated solution of calcium sulfate in sulfuric  
7 acid having the same acid normality, and wherein the  
8 AGIIS is non-volatile at room temperature and pressure.

52.       The method of claim 50, wherein the first  
2 tissue and the second tissue are animal tissues or plant  
3 tissues.

53.       A method for disinfecting a tissue, comprising:  
2           contacting the tissue with AGIIS.

54.       The method of claim 53, wherein the AGIIS is  
2 prepared by mixing calcium hydroxide with sulfuric acid  
3 with or without calcium sulfate added thereto, and the  
4 AGIIS having a certain acid normality is less effective  
5 in charring sucrose and less corrosive to an animal skin  
6 than a saturated solution of calcium sulfate in sulfuric  
7 acid having the same acid normality, and wherein the  
8 AGIIS is non-volatile at room temperature and pressure.  
9

-65-

55. The method of claim 53, wherein the tissue is  
2 an animal tissue or a plant tissue.

56. A method for cleaning a product, comprising:  
2 contacting the product with AGIIS.

57. The method of claim 56, wherein the AGIIS is  
2 prepared by mixing calcium hydroxide with sulfuric acid  
3 with or without calcium sulfate added thereto, and the  
4 AGIIS having a certain acid normality is less effective  
5 in charring sucrose and less corrosive to an animal skin  
6 than a saturated solution of calcium sulfate in sulfuric  
7 acid having the same acid normality, and wherein the  
8 AGIIS is non-volatile at room temperature and pressure.

58. The method of claim 56, wherein the product is  
2 a tissue, a microelectronic product, or a construction  
3 product.

59. The method of claim 58, wherein the  
2 construction product is new or reworked.

60. A method for synchronizing a harvest of a  
2 desired part of a plant, comprising:  
3 contacting the desired part of the plant with AGIIS.

61. The method of claim 60, wherein the AGIIS is  
2 prepared by mixing calcium hydroxide with sulfuric acid

-66-

3 with or without calcium sulfate added thereto, and the  
4 AGIIS having a certain acid normality is less effective  
5 in charring sucrose and less corrosive to an animal skin  
6 than a saturated solution of calcium sulfate in sulfuric  
7 acid having the same acid normality, and wherein the  
8 AGIIS is non-volatile at room temperature and pressure.  
9

62. A method for preserving or improving  
2 organoleptic quality of a desired part of a plant,  
3 comprising:  
4 contacting the desired part of the plant with AGIIS.

63. The method of claim 62, wherein the AGIIS is  
2 prepared by mixing calcium hydroxide with sulfuric acid  
3 with or without calcium sulfate added thereto, and the  
4 AGIIS having a certain acid normality is less effective  
5 in charring sucrose and less corrosive to an animal skin  
6 than a saturated solution of calcium sulfate in sulfuric  
7 acid having the same acid normality, and wherein the  
8 AGIIS is non-volatile at room temperature and pressure.

64. The method of claim 62, wherein the contacting  
2 of the desired part of the plant occurs prior to harvest,  
3 during harvest, during handling, or post harvest.

65. A method for reducing biological contaminants  
2 in water, comprising:

-67-

3 adding to the water a sufficient amount of AGIIS to  
4 reduce the biological contaminants.

66. The method of claim 65, wherein the AGIIS is  
2 prepared by mixing calcium hydroxide with sulfuric acid  
3 with or without calcium sulfate added thereto, and the  
4 AGIIS having a certain acid normality is less effective  
5 in charring sucrose and less corrosive to an animal skin  
6 than a saturated solution of calcium sulfate in sulfuric  
7 acid having the same acid normality, and wherein the  
8 AGIIS is non-volatile at room temperature and pressure.

67. The method of claim 65, wherein the water is  
2 portable water, storm sewer water, or sanitary sewer  
3 water.

68. A method for preparing AGIIS comprising:  
2 preparing an aqueous solution of a mineral acid;  
3 preparing an aqueous solution or slurry of a Group  
4 IIA hydroxide or a Group IIA salt;  
5 mixing the aqueous solution of the mineral acid with  
6 the aqueous solution or slurry of the Group IIA hydroxide  
7 or the Group IIA salt;  
8 removing solid formed to isolate AGIIS, and the  
9 AGIIS having a certain acid normality is less effective  
10 in charring sucrose and less corrosive to an animal skin  
11 than a saturated solution of calcium sulfate in sulfuric  
12 acid having the same acid normality, and wherein the  
13 AGIIS is non-volatile at room temperature and pressure.

-68-

69. A method for preparing AGIIS comprising:

2 mixing a mineral acid in water with a Group IIA  
3 hydroxide and the resultant AGIIS having a certain acid  
4 normality is less effective in charring sucrose and less  
5 corrosive to an animal skin than a saturated solution of  
6 calcium sulfate in sulfuric acid having the same acid  
7 normality, and wherein the AGIIS is non-volatile at room  
8 temperature and pressure.

70. The method of claim 69, wherein the mineral

2 acid is sulfuric acid and the Group IIA hydroxide is  
3 calcium hydroxide.

71. The method of claim 70, wherein for every mole

2 of sulfuric acid used, the amount of calcium hydroxide  
3 ranges from about 0.1 mole to about 0.5 mole.

72. A method of preparing AGIIS comprising:

2 adding a predetermined amount of calcium sulfate to  
3 an aqueous solution of concentrated sulfuric acid to give  
4 a mixture;

5 adding a calculated amount of slurry of calcium  
6 hydroxide in water to the mixture to give a reacted  
7 mixture;

8 removing solid formed in the reacted mixture to give  
9 the AGIIS; and the AGIIS having a certain acid  
10 normality is less effective in charring sucrose and less  
11 corrosive to an animal skin than a saturated solution of

-69-

12 calcium sulfate in sulfuric acid having the same acid  
13 normality, and wherein the AGIIS is non-volatile at room  
14 temperature and pressure.

73. The method of claim 72, further comprising  
2 introducing gaseous carbon dioxide into the mixture of  
3 the sulfuric acid containing calcium sulfate and the  
4 calcium hydroxide.

74. The method of claim 70, wherein for every mole  
2 of sulfuric acid used, the amount of calcium hydroxide  
3 ranges from about 0.1 mole to about 0.5 mole.

75. A method for preparing AGIIS having a desired  
2 final acid normality, comprising:  
3 (a) determining the amount of a mineral acid needed  
4 by the following equation:

$$5 \quad E_1 = (N/2) + (N/2 + B)$$

6 wherein  $E_1$  is the amount of the mineral  
7 acid, in moles, required before making purity adjustment;  
8  $N$  is the desired final acid normality; and  $B$  is the mole  
9 ratio of a Group IIA hydroxide to the mineral acid needed  
10 to obtain the AGIIS having  $N$ , and  $B$  is derived from a  
11 pre-plotted curve depicting the relationship of the  
12 mineral acid and the Group IIA hydroxide for a desired  $N$ ;

13 (b) making purity adjustment for the mineral acid  
14 used by the following equation:

$$15 \quad E_2 = E_1/C$$

-70-

16                wherein  $E_2$  is the amount of the mineral acid, in  
17 moles, required after purity adjustment;  $E_1$  is as defined  
18 above; and C is the purity adjustment factor for the  
19 mineral acid;

20                (c) determining the amount of water, in ml, needed  
21 to be added to the mineral acid by the following  
22 equation:

$$23 \qquad \qquad \qquad G = J - E_2 - I$$

24                wherein G is the amount of water, in ml,  
25 required to be added to the mineral acid; J is the final  
26 volume of aqueous mineral acid solution; I is the volume  
27 amount of Group IIA hydroxide needed, given below; and  $E_2$   
28 is as defined above;

29                (d) adding G to  $E_2$  to give the final aqueous  
30 solution of the mineral acid, wherein both G and  $E_2$  are as  
31 defined above;

32                (e) determining the amount of Group IIA hydroxide,  
33 in moles, needed by the following equation:

$$34 \qquad \qquad \qquad F_1 = N/2 \times B$$

35                wherein  $F_1$  is the amount of Group IIA hydroxide,  
36 in moles, needed before making purity adjustment; and B  
37 and N are as defined above;

38                (f) making purity adjustment for the Group IIA  
39 hydroxide used by the following equation:

$$40 \qquad \qquad \qquad F_2 = F_1/D$$

41                wherein  $F_2$  is the amount of the Group IIA  
42 hydroxide, in moles, required after purity adjustment;  $F_1$   
43 is as defined above; and D is the purity adjustment  
44 factor for the Group IIA hydroxide;

-71-

45 (g) determining the amount of water, in ml, needed  
46 to make the solution or slurry of Group IIA hydroxide by  
47 the following equation:

48 
$$H = F_2 \times 1.5$$
 wherein H is the amount of  
49 water, in ml, needed to make the solution or slurry of  
50 Group IIA hydroxide; and  $F_2$  is as defined above;

51 (h) determining the amount of the aqueous solution  
52 or slurry of Group IIA hydroxide, in ml, needed to be  
53 added to the aqueous solution of mineral acid to give the  
54 AGIIS with a desired final acid normality by the  
55 following equation:

56 
$$I = F_2 \times 2$$

57 wherein I is the amount of Group IIA hydroxide  
58 solution or slurry, in ml, needed; and  $F_2$  is as defined  
59 above;

60 (i) adding H to  $F_2$  to give the final aqueous  
61 solution or slurry of Group IIA hydroxide, wherein both H  
62 and  $F_2$  are as defined above;

63 (j) adding the final aqueous solution or slurry of  
64 Group IIA hydroxide of (i) to the final aqueous solution  
65 of mineral acid of (d);

66 (k) allowing the final aqueous solution or slurry  
67 of Group IIA hydroxide and the final aqueous solution of  
68 mineral acid of (j) to react; and

69 (l) removing solid formed from (K).

76. The method of claim 75 further comprising  
2 adding a Group IIA salt of a dibasic acid to the final  
3 aqueous mineral acid solution of (d).

-72-

77. The method of claim 76, wherein the mineral  
2 acid is sulfuric acid, the Group IIA hydroxide is calcium  
3 hydroxide, and the Group IIA salt of a dibasic acid is  
4 calcium sulfate.

78. The AGIIS having a desired final acid normality  
2 prepared by the method of claim 75.




Fig. 1



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
24 August 2000 (24.08.2000)

PCT

(10) International Publication Number  
**WO 00/48477 A3**

(51) International Patent Classification<sup>7</sup>: **A23L 3/358**,  
1/304, A61K 33/08, 33/06, C01F 11/46 // (A61K 33/08,  
33:06, 33:04)

CA 95678 (US). YAO, Yu [US/US]; 401 Rocky Point  
Drive #5, Roseville, CA 95678 (US). LEWIS, David, E.  
[US/US]; 816 3rd Avenue, Eau Claire, WI 54703 (US).

(21) International Application Number: PCT/US00/03993

(74) Agent: CHWANG, T., Ling; Hitt Chwang & Gaines,  
P.C., Suite 225, 275 West Campbell Road, Richardson, TX  
75080 (US).

(22) International Filing Date: 14 February 2000 (14.02.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
09/253,482 19 February 1999 (19.02.1999) US  
Not furnished 9 February 2000 (09.02.2000) US

(71) Applicant (for all designated States except US): MORN-  
INGSTAR DIAGNOSTICS, INC. [US/US]; Suite 130,  
1376 Lead Hill Boulevard, Roseville, CA 95661 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): KEMP, Maurice,  
Clarence [US/US]; 4401 Woodthrush Drive, El Dorado  
Hills, CA 95762 (US). LALUM, Robert, Blaine, Lalum  
[US/US]; 6949 Greenback Lane #B, Citrus Heights, CA  
95621 (US). XIE, Zhong, Wei [US/US]; 1141 Elderberry  
Circle, Folsom, CA 95630 (US). CUNHA, Michael, An-  
thony [US/US]; 355 Shadow Ridge Drive #3, Roseville,  
CA 95678 (US). CARPENTER, Robert, H. [US/US];  
1303 Pecan Street, Bastrop, TX 78602 (US). ZHANG,  
Shu [US/US]; 401 Rocky Point Drive #5, Roseville,

(81) Designated States (national): AE, AL, AM, AT, AU, AZ,  
BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK,  
DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL,  
IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU,  
LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT,  
RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA,  
UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM,  
KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent  
(AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent  
(AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU,  
MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM,  
GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— With international search report.

(88) Date of publication of the international search report:  
30 November 2000

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

(54) Title: ACIDIC SOLUTION OF SPARINGLY-SOLUBLE GROUP IIA COMPLEXES

(57) Abstract: An acidic solution of sparingly-soluble Group IIA complexes ("AGIIS"), its preparation and its uses. The AGIIS can be prepared by mixing a mineral acid (such as sulfuric acid), and a Group IIA hydroxide (such as calcium hydroxide) or a Group IIA salt of a dibasic acid (such as calcium sulfate), or a mixture of the two Group IIA compounds, followed by removing the solid formed. The various uses include cleaning, food production, decontamination, bioremediation, agricultural application, medical application, and detoxification of substances.

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/03993

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A23L3/358 A23L1/304 A61K33/08 A61K33/06 C01F11/46  
 //(A61K33/08,33:06,33:04)

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A23L A61K A23B C01F

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

WPI Data, PAJ, FSTA

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5 264 017 A (VAN DE WALLE RICHARD H) 23 November 1993 (1993-11-23)  column 1, line 11-19 column 2, line 60 -column 3, line 14 claims 1,12	1,2,6, 12,15, 19,22, 27,29, 30,32, 36,69
P,X	WO 99 23896 A (NESTLE SA ;MALLANGI CHANDRASEKHARA REDDY (US); SHER ALEXANDER (US)) 20 May 1999 (1999-05-20) page 3, line 32 -page 4, line 6 page 4, line 29-34 page 5, line 19-33 page 6, line 19-25 claims 1,2,5,6,11-16	1,2,6, 12,15, 68,69
	-/-	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

14 June 2000

Date of mailing of the international search report

28. 08. 2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
 NL - 2280 HV Rijswijk  
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
 Fax (+31-70) 340-3016

Authorized officer

Piret-Viprey, E

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/03993

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5 830 838 A (OVERTON JAMES M ET AL) 3 November 1998 (1998-11-03) cited in the application  column 1, line 10-15 column 1, line 52 -column 2, line 26 claim 1	24-26, 30-32, 56-59, 68-71,78
X	US 5 571 336 A (WURZBURGER STEPHEN R ET AL) 5 November 1996 (1996-11-05)  figure I column 2, line 1-15 claim 1	24-26, 30-32, 56-59, 68-70,78
X	US 4 306 516 A (CURREY JOHN E) 22 December 1981 (1981-12-22) column 2, line 14-30 column 3, line 60 -column 4, line 14 claims 1-10	14,30,32
X	DATABASE WPI Derwent Publications Ltd., London, GB; AN 1981-54598 XP002139494 & SU 778 753 A (TOPCHIEV PETROCHEM SYNTH), 15 November 1980 (1980-11-15) abstract	69,70
A		72,78
X	DATABASE WPI Derwent Publications Ltd., London, GB; AN 1989-003849 XP002140034 & JP 63 282117 A (SHIMIZU N.), 18 November 1988 (1988-11-18) abstract	68-70,78
A		75
X	GB 1 545 788 A (ICI LTD) 16 May 1979 (1979-05-16) example 1 claims 1,13-15	68,69
X	WO 94 09798 A (BONAPARTE CO) 11 May 1994 (1994-05-11)  page 5, line 21 -page 6, line 14 page 10, line 29 -page 11, line 5 claims 1,2,9,37-45	1,2,6, 12,13, 53,55
	-/-	

# INTERNATIONAL SEARCH REPORT

Inter. Application No

PCT/US 00/03993

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>WO 96 01566 A (DARATECH PTY LTD ;REYES VICENTE GERONIMO (AU)) 25 January 1996 (1996-01-25)</p> <p>page 1, line 1-9 page 4, line 6 -page 5, line 21 claims 1,2,8,9,18,22,23,25,32,34</p>	<p>1,2,6, 12,13, 15,27, 62,64</p>

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US 00/03993

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:  
  
Although claims 42-49 are directed to a diagnostic method practised on the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☒ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:  
  
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

## FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

## Continuation of Box I.2

Present claims 1,2 relate to an extremely large number of possible compounds. Support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT is to be found, however, for only a very small proportion of the compounds claimed. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Consequently, the search has been carried out for those parts of the claims which appear to be supported and disclosed, namely those parts relating to the compounds:

## - claim 3:

mineral acid= sulfuric acid

Group IIA hydroxide= calcium hydroxide

Group IIA salt of a dibasic acid= calcium sulfate

## - description: pagell, line 22 - page 12, line 20

mineral acid= sulfuric acid

Group IIA hydroxide= calcium hydroxide

Group IIA salt of a dibasic acid= calcium sulfate, calcium oxide, calcium carbonate.

## - examples: 1-3

mineral acid= sulfuric acid

Group IIA hydroxide= calcium hydroxide

Group IIA salt of a dibasic acid= calcium sulfate.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 00/03993

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 5264017 A	23-11-1993	CA 1318147 A	25-05-1993
WO 9923896 A	20-05-1999	US 6024994 A AU 1152899 A	15-02-2000 31-05-1999
US 5830838 A	03-11-1998	US 5895782 A	20-04-1999
US 5571336 A	05-11-1996	NONE	
US 4306516 A	22-12-1981	NONE	
SU 778753 A	15-11-1980	NONE	
JP 63282117 A	18-11-1988	NONE	
GB 1545788 A	16-05-1979	NONE	
WO 9409798 A	11-05-1994	AU 5589594 A CA 2148256 A EP 0666750 A JP 8502976 T	24-05-1994 11-05-1994 16-08-1995 02-04-1996
WO 9601566 A	25-01-1996	AU 2916895 A	09-02-1996



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

CORRECTED VERSION

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
24 August 2000 (24.08.2000)

PCT

(10) International Publication Number  
**WO 00/48477 A3**

(51) International Patent Classification: **A23L 3/358**,  
1/304, A61K 33/08, 33/06, C01F 11/46 // (A61K 33/08,  
33/06, 33/04)

(74) Agent: CHWANG, T., Ling; Jackson Walker, LLP, Suite  
600, 2435 North Central Expressway, Richardson, TX  
75080 (US).

(21) International Application Number: PCT/US00/03993

(22) International Filing Date: 14 February 2000 (14.02.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
(9/253,482 19 February 1999 (19.02.1999) US  
(9/500,473 9 February 2000 (09.02.2000) US

(71) Applicant (for all designated States except US): MORN-  
INGSTAR DIAGNOSTICS, INC. [US/US]; Suite 130,  
1376 Lead Hill Boulevard, Roseville, CA 95661 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): KEMP, Maurice,  
Clarence [US/US]; 4401 Woodthrush Drive, El Dorado  
Hills, CA 95762 (US). LALUM, Robert, Blaine, Lalum  
[US/US]; 6949 Greenback Lane #B, Citrus Heights, CA  
95621 (US). XIE, Zhong, Wei [US/US]; 1141 Elderberry  
Circle, Folsom, CA 95630 (US). CUNHA, Michael, An-  
thony [US/US]; 355 Shadow Ridge Drive #3, Roseville,  
CA 95678 (US). CARPENTER, Robert, H. [US/US];  
1303 Pecan Street, Bastrop, TX 78602 (US). ZHANG,  
Shu [US/US]; 401 Rocky Point Drive #5, Roseville,  
CA 95678 (US). YAO, Yu [US/US]; 401 Rocky Point  
Drive #5, Roseville, CA 95678 (US). LEWIS, David, E.  
[US/US]; 816 3rd Avenue, Eau Claire, WI 54703 (US).

(81) Designated States (national): AE, AL, AM, AT, AU, AZ,  
BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK,  
DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL,  
IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU,  
LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT,  
RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA,  
UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM,  
KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent  
(AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent  
(AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU,  
MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM,  
GA, GN, GW, ML, MR, NE, SN, TD, TG).

**Published:**

- With international search report.
- With amended claims and statement.

(88) Date of publication of the international search report:  
30 November 2000

(48) Date of publication of this corrected version:  
3 May 2001

(15) Information about Correction:  
see PCT Gazette No. 18/2001 of 3 May 2001, Section II

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

(54) Title: ACIDIC SOLUTION OF SPARINGLY-SOLUBLE GROUP IIA COMPLEXES

(57) Abstract: An acidic solution of sparingly-soluble Group IIA complexes ("AGIIS"), its preparation and its uses. The AGIIS can be prepared by mixing a mineral acid (such as sulfuric acid), and a Group IIA hydroxide (such as calcium hydroxide) or a Group IIA salt of a dibasic acid (such as calcium sulfate), or a mixture of the two Group IIA compounds, followed by removing the solid formed. The various uses include cleaning, food production, decontamination, bioremediation, agricultural application, medical application, and detoxification of substances.

WO 00/48477 A3

